

CHOW MOTIVE OF FULTON-MACPHERSON CONFIGURATION SPACES AND WONDERFUL COMPACTIFICATIONS

LI LI

1. INTRODUCTION

The purpose of this article is to study the Chow groups and Chow motives of the so-called wonderful compactifications of an arrangement of subvarieties, in particular the Fulton-MacPherson configuration spaces.

All the varieties in the paper are over an algebraically closed field. Let Y be a nonsingular quasi-projective variety. Let $\mathcal{S}$ be an arrangement of subvarieties of Y (cf. Definition 2.2). Let $\mathcal{G}$ be a building set of $\mathcal{S}$, i.e., a finite set of nonsingular subvarieties in $\mathcal{S}$ satisfying Definition 2.3. The wonderful compactification $Y_{\mathcal{G}}$ is constructed by blowing up Y along subvarieties in $\mathcal{G}$ successively (cf. Definition 2.5). There are different orders in which the blow-ups can be carried out, for example we can blow up along the centers in any order that is compatible with the inclusion relation. There are many important examples of such compactifications: De Concini and Procesi's wonderful model of a subspace arrangement, the Fulton-MacPherson configuration spaces, the moduli space $\overline{\mathcal{M}}_{0,n}$ of stable rational curves with n marked points, etc. These spaces have many properties in common. Studying them by a uniform method gives us better understanding of these spaces. In this article, we study their Chow groups and Chow motives.

If we assume that Y is projective, then the Chow motive of $Y_{\mathcal{G}}$, denoted by $h(Y_{\mathcal{G}})$, can be decomposed canonically into a direct sum of the motive of Y and the twisted motives of the subvarieties in the arrangement (cf. §2.1 for a review of Chow motives). We will prove the following theorem, where the precise definition of the set $M_{\mathcal{T}}$ and the subvarieties $Y_0\mathcal{T}$ of Y are in §3.

Main Theorem (Theorems 3.1, 3.2). *Let Y be a nonsingular quasi-projective variety, $\mathcal{G}$ be a building set and $Y_{\mathcal{G}}$ be the wonderful compactification $Y_{\mathcal{G}}$. Then we have the Chow group decomposition*

$$A^*Y_{\mathcal{G}} = A^*Y \oplus \bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}} A^{*-||\underline{\mu}||}(Y_0\mathcal{T})$$

2000 *Mathematics Subject Classification.* 14C15, 14N20.

where $\mathcal{T}$ runs through all $\mathcal{G}$ -nests. Moreover, when Y is projective we have a Chow motive decomposition

$$h(Y_{\mathcal{G}}) \cong h(Y) \oplus \bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}} h(Y_0 \mathcal{T})(\|\underline{\mu}\|)$$

where $\mathcal{T}$ runs through all $\mathcal{G}$ -nests. In this case the correspondences giving the above isomorphism are canonical in the following sense: although there is no canonical order of blow-ups (in general) to construct $Y_{\mathcal{G}}$, the correspondences turn out to be independent of the order we choose.

The Fulton-MacPherson configuration space $X[n]$ is one of the most interesting examples of the wonderful compactification $Y_{\mathcal{G}}$ where $Y = X^n$ and $\mathcal{G}$ is the set of all the diagonals in X^n (cf. §4.1). Applying the main theorem to $X[n]$, we obtain the following theorem, where the precise definition of the nests $\mathcal{S}$, the polydiagonals $\Delta_{\mathcal{S}}$, the integers $c(\mathcal{S})$, the sets of lattice points $M_{\mathcal{S}}$, and the correspondences $\alpha_{\mathcal{S}, \underline{\mu}}$ and $\beta_{\mathcal{S}, \underline{\mu}}$ are in §4.1.

Theorem 4.2. *Let X be a nonsingular projective variety. Then there is a canonical isomorphism of Chow motives*

$$\bigoplus_{\mathcal{S}} \bigoplus_{\underline{\mu} \in M_{\mathcal{S}}} \alpha_{\mathcal{S}, \underline{\mu}} : h(X[n]) \cong \bigoplus_{\mathcal{S}} \bigoplus_{\underline{\mu} \in M_{\mathcal{S}}} h(\Delta_{\mathcal{S}})(\|\underline{\mu}\|)$$

with the inverse $\sum_{\mathcal{S}} \sum_{\underline{\mu} \in M_{\mathcal{S}}} \beta_{\mathcal{S}, \underline{\mu}}$. Equivalently, we have the following decomposition of the Chow motive of $X[n]$:

$$h(X[n]) \cong \bigoplus_{\mathcal{S}} \bigoplus_{\underline{\mu} \in M_{\mathcal{S}}} h(X^{c(\mathcal{S})})(\|\underline{\mu}\|).$$

Here are two consequences of this theorem. One is that we can easily express the decomposition of $h(X[n])$ using a generating function $N(x, t)$, as follows.

Theorem 4.3. *Define $f_i(x)$ to be the polynomials whose exponential generating function $N(x, t) = \sum_{i \geq 1} f_i(x) \frac{t^i}{i!}$ satisfies the identity*

$$(1-x)x^d t + (1-x^{d+1}) = \exp(x^d N) - x^{d+1} \exp(N)$$

where $d = \dim X$. Then

$$h(X[n]) = \bigoplus_{\substack{1 \leq k \leq n \\ i \geq 0}} (h(X^k)(i))^{\oplus [\frac{x^i t^n}{n!}] \frac{N^k}{k!}}.$$

The other consequence is a decomposition of the Chow motive of the quotient variety $X[n]/\mathfrak{S}_n$ obtained from the natural symmetric group $\mathfrak{S}_n$ action on $X[n]$. To make sense of the motive of a quotient variety, we assume the base field is of characteristic 0. The correspondences appeared in Theorem 4.2 are canonical, and therefore symmetric with respect to the

symmetric group $\mathfrak{S}_n$. It is then possible to compute the $\mathfrak{S}_n$ -invariant part of $h(X[n])$, which is the Chow motive of $X[n]/\mathfrak{S}_n$. As pointed out by [FM94], unlike the isotropy groups of a point in X^n , the isotropy group of any point in $X[n]$ is always solvable, therefore the singularity of $X[n]/\mathfrak{S}_n$ is “better” than the singularity of the symmetric product $X^{(n)} := X^n/\mathfrak{S}_n$. It would be interesting to see how different is the Chow motive $h(X[n]/\mathfrak{S}_n)$ from $h(X^{(n)})$. In the following theorem, an unlabeled weighted forest is a forest whose nodes are not labeled and that each non-leaf node is attached by a positive integer called weight; we call an unlabeled weighted forest of type $\nu := \{n_1, \dots, n_r\}$ if the forest is of the form $n_1 T_1 + \dots + n_r T_r$, where T_i are mutually distinct unlabeled weighted tree.

Theorem 5.3. *For any unordered set of positive integers $\nu = \{n_1, \dots, n_r\}$ and any non-negative integer m , let $\lambda(\nu, m)$ to be the number of unlabeled weighted forest with n leaves, of type ν and of total weight m , such that at each non-leaf v with c_v children, the weight m_v satisfies $1 \leq m_v \leq (c_v - 1) \dim X - 1$. Then*

$$h(X[n]/\mathfrak{S}_n) = \bigoplus_{\nu, m} \left[h(X^{(n_1)} \times \dots \times X^{(n_r)})(m) \right]^{\oplus \lambda(\nu, m)}.$$

The importance of all the above results of Chow motives can be seen through a working principle:

Principle: *A result proved for Chow motives is valid if we replace them by homological/numerical motives, Chow groups $A_{\mathbb{Q}}^*$, cohomology groups $H_{\mathbb{Q}}^*$, Grothendieck groups (the aforementioned groups are taken with $\mathbb{Q}$ -coefficients), Hodge structures, etc.*

Thus for example, we have a decomposition for the $\mathbb{Q}$ -coefficient singular cohomology of $Y_{\mathcal{G}}$, $X[n]$ and $X[n]/\mathfrak{S}_n$.

The paper is organized as follows. §2 contains a review of motives and the wonderful compactifications of arrangement of subvarieties. In §3 a motivic decomposition for the wonderful compactifications is proved. In §4 we give a motivic decomposition for the Fulton-MacPherson configuration spaces. §5 gives a motivic decomposition for the quotient variety $X[n]/\mathfrak{S}_n$.

Acknowledgements. The paper is based on part of the author’s thesis. In many ways I am greatly indebted to Mark de Cataldo, my Ph.D. advisor. I would also thank Blaine Lawson, Sorin Popescu, Dror Varolin, Byungheup Jun for encouragement and useful discussions. The author thanks the referee for the detailed review of the first version with many very helpful suggestions to clarify and simplify the paper.

2. PRELIMINARIES

2.1. Motives. Given an algebraic variety X of dimension d , let $A^i X = A_{d-i} X$ be the Chow group of codimension i , i.e., the group of algebraic

cycles of codimension i in X modulo rational equivalence. Define $A_{\mathbb{Q}}^i X = A^i X \otimes_{\mathbb{Z}} \mathbb{Q}$.

Let X, Y be two non-singular projective varieties. The group of *correspondences of degree r* from X to Y is defined as

$$\text{Corr}^r(X, Y) := A^{\dim X + r}(X \times Y).$$

The group $\text{Corr}_{\mathbb{Q}}^r(X, Y)$ denotes the tensor of $\text{Corr}^r(X, Y)$ with $\mathbb{Q}$.

The *composition* of two correspondences $f \in \text{Corr}^r(X_1, X_2)$ and $g \in \text{Corr}^s(X_2, X_3)$ is a correspondence in $\text{Corr}^{r+s}(X_1, X_3)$ defined as

$$g \circ f := \pi_{13*}(\pi_{12}^* f \cdot \pi_{23}^* g)$$

where π_{ij} is the projection from $X_1 \times X_2 \times X_3$ to $X_i \times X_j$.

A correspondence $p \in \text{Corr}^0(X, X)$ is called a *projector* of X if $p^2 := p \circ p = p$.

Let $\mathcal{V}$ denote the category of (not necessarily connected) non-singular projective varieties over a field k .

Definition 2.1 ([CH00]). *The category of Chow motives over k , denoted by CHM , is defined as follows: an object of CHM , called a Chow motive, is a triple (X, p, r) , where X is a nonsingular projective variety, p is a projector of X and r is an integer. The morphisms in CHM are defined as*

$$\text{Hom}_{CHM}((X, p, r), (Y, q, s)) := q \circ \text{Corr}^{s-r}(X, Y) \circ p.$$

The composition of morphisms is defined as the composition of correspondences.

For a Chow motive $M = (X, p, r)$ and an integer ℓ , we define

$$M(\ell) := (X, p, r + \ell).$$

There is a natural contravariant functor h from $\mathcal{V}$ to CHM , which sends X to $(X, id_X, 0)$ and sends a morphism $f : X \rightarrow Y$ to $\Gamma_f^t : h(Y) \rightarrow h(X)$, the transpose of the graph of f . Naturally, $h(X)(\ell)$ stands for the Chow motive (X, id_X, ℓ) .

According to [dBV98], we can generalize the theory of Chow motives on nonsingular projective varieties to the one on varieties which are quotients of smooth projective varieties by finite group actions. To be more precisely, let $\mathcal{V}'$ be the category of varieties of type X/G with $X \in Ob\mathcal{V}$ and G a finite group. We can define the group of correspondences $\text{Corr}_{\mathbb{Q}}^r(X', Y')$ for $X', Y' \in \mathcal{V}'$ and the category of Chow motives CHM' similar to the nonsingular case. (The difference is that we have to use $\mathbb{Q}$ -coefficients). There is a natural contravariant functor $h : \mathcal{V}' \rightarrow CHM'$.

Define the G -average correspondence ave_G as

$$\text{ave}_G := \frac{1}{|G|} \sum [g] \in \text{Corr}_{\mathbb{Q}}^0(X, X)$$

where $[g]$ is given by the graph of g in $X \times X$. By [dBV98] Proposition 1.2, there is an isomorphism

$$h(X/G) \cong (X, \text{ave}\Delta) \cong h(X)^G.$$

Such a definition is consistent with the realization functors and $\mathbb{Q}$ -coefficient Chow groups.

2.2. Wonderful compactification of an arrangement of subvarieties.

The wonderful compactification of an arrangement of subvarieties is introduced in [Li06] as a generalization of De Concini and Procesi's wonderful model of subspace arrangements. We briefly review the definition and some properties of such compactifications. For details we refer to [Li06].

Definition 2.2. *A (simple) arrangement of subvarieties of Y is a finite set $\mathcal{S} = \{S_i\}$ of nonsingular closed subvarieties of Y satisfying the following conditions:*

- (1) S_i and S_j intersect cleanly (i.e. their intersection is nonsingular and $T(S_i \cap S_j) = T(S_i)|_{(S_i \cap S_j)} \cap T(S_j)|_{(S_i \cap S_j)}$),
- (2) $S_i \cap S_j$ is either empty or equal to some $S_k \in \mathcal{S}$.

Definition 2.3. *Let $\mathcal{S}$ be an arrangement of subvarieties of Y . A subset $\mathcal{G} \subseteq \mathcal{S}$ is called a building set of $\mathcal{S}$ if $\forall S \in \mathcal{S}$, the minimal elements in $\mathcal{G}$ which contains S intersect transversally and their intersection is S (this condition is always satisfied if $S \in \mathcal{G}$). These minimal elements are called the $\mathcal{G}$ -factors of S . We call a finite set $\mathcal{G}$ of subvarieties a building set if the set*

$$\mathcal{S} := \left\{ \bigcap_{V \in \mathcal{F}} V \right\}_{\mathcal{F}},$$

where $\mathcal{F}$ runs through all subsets of $\mathcal{G}$, is an arrangement and $\mathcal{G}$ is a building set of $\mathcal{S}$ (for $\mathcal{F} = \emptyset$ we set $\bigcap_{V \in \mathcal{F}} V = Y$). In this case we call $\mathcal{S}$ the induced arrangement of $\mathcal{G}$.

Definition 2.4. *Let $\mathcal{G}$ be a building set. A subset $\mathcal{T} \subseteq \mathcal{G}$ is called $\mathcal{G}$ -nested (or a $\mathcal{G}$ -nest) if it satisfies one of the following equivalent relations:*

- (1) *There is a flag of elements in $\mathcal{S}$: $S_1 \subseteq S_2 \subseteq \cdots \subseteq S_k$, such that*

$$\mathcal{T} = \bigcup_{i=1}^k \{A : A \text{ is a } \mathcal{G}\text{-factor of } S_i\}.$$

(We say that $\mathcal{T}$ is induced by the flag $S_1 \subseteq S_2 \subseteq \cdots \subseteq S_k$.)

- (2) *Let $A_1, \dots, A_k$ be the minimal elements of $\mathcal{T}$, then they are all the $\mathcal{G}$ -factors of a certain element in $\mathcal{S}$, and for each $1 \leq i \leq k$, the set $\{A \in \mathcal{T} : A \supsetneq A_i\}$ is also $\mathcal{G}$ -nested defined by induction.*

The wonderful compactification is defined as follows:

Definition 2.5. Denote $Y^\circ = Y \setminus \bigcup_{G \in \mathcal{G}} G$. There is a natural locally closed embedding

$$Y^\circ \hookrightarrow Y \times \prod_{G \in \mathcal{G}} \text{Bl}_G Y.$$

The closure of this embedding, denoted by $Y_{\mathcal{G}}$, is called the wonderful compactification of $\mathcal{G}$.

The wonderful compactification $Y_{\mathcal{G}}$ of $\mathcal{G}$ has the following properties, where (1) and (2) are in Theorem 1.2 in [Li06] and (3) is clear from the proof there.

Theorem 2.6. The variety $Y_{\mathcal{G}}$ is nonsingular. For each $G \in \mathcal{G}$ there is a nonsingular divisor D_G on $Y_{\mathcal{G}}$ such that:

- (1) The union of the divisors D_G is $Y_{\mathcal{G}} \setminus Y^\circ$,
- (2) Any collection of the divisors D_G intersects transversally. An intersection of divisors $D_{T_1} \cap \cdots \cap D_{T_r}$ is nonempty exactly when $\{T_1, \dots, T_r\}$ forms a $\mathcal{G}$ -nest.
- (3) Each D_G is the unique connected component of $\pi^{-1}(G)$ that maps surjectively to the subvariety G , where π is the natural morphism $Y_{\mathcal{G}} \rightarrow Y$. (This D_G is called the dominant transform of G and denoted by $\tilde{G}$ in [Li06].)

The dominant transform can also be defined as follows. Let $\pi : \tilde{Y} \rightarrow Y$ be the blow-up along a nonsingular subvariety $G \subsetneq Y$. For any irreducible subvariety V in Y , we define the dominant transform of V , denoted by $\tilde{V}$ or $V^\sim$, to be the strict transform of V when $V \not\subseteq G$, and $\pi^{-1}(V)$ when $V \subseteq G$. For a sequence of N blow-ups $Y_N \rightarrow Y_{N-1} \rightarrow \cdots \rightarrow Y_1 \rightarrow Y_0$ and a subvariety $V \subseteq Y_0$ we define the dominant transform $\tilde{V} \subseteq Y_N$ (or denoted by $V^\sim$) to be the N -th iterated dominant transform $(\cdots ((V^\sim)^\sim) \cdots)^\sim$.

It is known (cf. [Li06]) that $Y_{\mathcal{G}}$ can be constructed by a sequence of blow-ups as follows. Let Y be a nonsingular variety, $\mathcal{S}$ be an arrangement of subvarieties and

$$\mathcal{G} = \{G_1, \dots, G_N\}$$

be a building set with respect to $\mathcal{S}$. Suppose the subvarieties in $\mathcal{G} = \{G_1, \dots, G_N\}$ are indexed in an order compatible with inclusion relations, i.e. $i \leq j$ if $G_i \subseteq G_j$. We define the triple $(Y_k, \mathcal{S}^{(k)}, \mathcal{G}^{(k)})$ inductively with respect to k , where Y_k is a nonsingular variety, $\mathcal{S}^{(k)}$ is an arrangement of subvarieties of Y_k and $\mathcal{G}^{(k)}$ is a building set with respect to $\mathcal{S}^{(k)}$:

(1) For $k = 0$, define $Y_0 = Y$, $\mathcal{S}^{(0)} = \mathcal{S}$, $\mathcal{G}^{(0)} = \mathcal{G} = \{G_1, \dots, G_N\}$, $G_i^{(0)} = G_i$ for $1 \leq i \leq N$.

(2) Assume the triple $(Y_k, \mathcal{S}^{(k)}, \mathcal{G}^{(k)})$ has been constructed. Define Y_{k+1} to be the blow-up of Y_k along the nonsingular subvariety $G_k^{(k)}$. Define $\mathcal{G}^{(k+1)}$ to be the dominant transform $(G^{(k)})^\sim$ for all $G \in \mathcal{G}$. Then $\mathcal{G}^{(k)} :=$

$\{G^{(k)}\}_{G \in \mathcal{G}}$ is a building set (by [Li06] Proposition 2.8). We denote the induced arrangement by $\mathcal{S}^{(k)}$.

(3) Continue the inductive construction until $k = N$. We get a nonsingular variety Y_N and all elements in the building set $\mathcal{G}^{(N)}$ are divisors. The resulting variety is isomorphic to $Y_{\mathcal{G}}$.

For any $\mathcal{G}$ -nest $\mathcal{T}$, define

$$Y_k \mathcal{T} = \bigcap_{G \in \mathcal{T}} G^{(k)}.$$

The following property of $Y_k \mathcal{T}$ is used often throughout the paper.

Proposition 2.7. *Let $0 \leq k \leq N - 2$ and let $\mathcal{T} \subseteq \{G_{k+2}, G_{k+3}, \dots, G_N\}$ be a $\mathcal{G}$ -nest. Then $Y_{k+1} \mathcal{T}$ is an irreducible nonsingular subvariety of Y_{k+1} with the following property:*

If $\{G_{k+1}\} \cup \mathcal{T}$ is not a $\mathcal{G}$ -nest, then $G_{k+1}^{(k)} \cap Y_k \mathcal{T} = \emptyset$ and $Y_{k+1} \mathcal{T} \cong Y_k \mathcal{T}$; otherwise, the intersection $G_{k+1}^{(k)} \cap Y_k \mathcal{T}$ is clean, $Y_{k+1} \mathcal{T}$ is isomorphic to the blow-up of $Y_k \mathcal{T}$ along $G_{k+1}^{(k)} \cap Y_k \mathcal{T}$ with exceptional divisor $G_{k+1}^{(k+1)} \cap Y_{k+1} \mathcal{T}$ (where the intersection is transverse), and the codimension of $G_{k+1}^{(k)} \cap Y_k \mathcal{T}$ in $Y_k \mathcal{T}$ equals to

$$\begin{cases} \dim \cap_{G_{k+1} \subsetneq G \in \mathcal{T}} G - \dim G_{k+1}, & \text{if } \{G : G_{k+1} \subsetneq G \in \mathcal{T}\} \neq \emptyset; \\ \dim Y - \dim G_{k+1}, & \text{otherwise.} \end{cases}$$

Proof. We prove the statement by induction on k . The case $k = 0$ is obvious. Now assume that the statement is true for k .

(i) Suppose that $\{G_{k+1}\} \cup \mathcal{T}$ is not a $\mathcal{G}$ -nest. We will show that $G_{k+1}^{(k)} \cap Y_k \mathcal{T} = \emptyset$. As a consequence we have $Y_{k+1} \mathcal{T} \cong Y_k \mathcal{T}$, since the center of the blow-up is away from $Y_k \mathcal{T}$.

We prove by contradiction. Assume $G_{k+1}^{(k)} \cap Y_k \mathcal{T} \neq \emptyset$. Since $\mathcal{T}$ is a $\mathcal{G}$ -nest, $\{G^{(k)}\}_{G \in \mathcal{T}}$ is a $\mathcal{G}^{(k)}$ -nest by [Li06] Proposition 2.8 (3). By Definition 2.4 (1), the nest $\{G^{(k)}\}_{G \in \mathcal{T}}$ is induced by a flag

$$S'_1 \subseteq S'_2 \subseteq \dots \subseteq S'_l$$

where $S'_i \in \mathcal{S}^{(k)}$. We claim that $\{G_{k+1}^{(k)}\} \cup \{G^{(k)}\}_{G \in \mathcal{T}} \subseteq \mathcal{G}^{(k)}$ is a $\mathcal{G}^{(k)}$ -nest induced by the flag

$$(1) \quad (G_{k+1}^{(k)} \cap S'_1) \subseteq S'_1 \subseteq S'_2 \subseteq \dots \subseteq S'_l.$$

Indeed, since $Y_k \mathcal{T} = S'_1$, we know $G_{k+1}^{(k)} \cap S'_1 \neq \emptyset$. By [Li06] Lemma 2.4

(ii), the $\mathcal{G}^{(k)}$ -factors of $G_{k+1}^{(k)} \cap S'_1$ are $G^{(k)}$ and some $\mathcal{G}^{(k)}$ -factors of S'_1 , hence our claim follows. Then [Li06] Proposition 2.8 (3) asserts that, since $\{G_{k+1}^{(k)}\} \cup \{G^{(k)}\}_{G \in \mathcal{T}}$ is a $\mathcal{G}^{(k)}$ -nest, $\{G_{k+1}\} \cup \mathcal{T}$ must be a $\mathcal{G}$ -nest. But by assumption $\{G_{k+1}\} \cup \mathcal{T}$ is not a $\mathcal{G}$ -nest, contradiction.

(ii) Suppose that $\mathcal{T} \cup \{G_{k+1}\}$ is a $\mathcal{G}$ -nest. Let the $\mathcal{G}^{(k)}$ -factors of $Y_k \mathcal{T}$ be $G'_1, \dots, G'_r$. Then they are minimal elements in the $\mathcal{G}^{(k)}$ -nest $\{G^{(k)}\}_{G \in \mathcal{T}}$, by

the definition of nest. Assume without loss of generality that the first m subvarieties $G'_1, \dots, G'_m$ contain $G_{k+1}^{(k)}$. Define $A = \cap_{i=1}^m G'_i$, $B = \cap_{i=m+1}^r G'_i$, then $Y_k \mathcal{T} = A \cap B$ is the $G_{k+1}^{(k)}$ -factorization of $Y_k \mathcal{T}$ by [Li06] Definition-Lemma 2.6.

Notice that for $p, q \geq k+2$ and $G_p^{(k)} \subseteq G_q^{(k)}$, we have $G_p^{(k+1)} \subseteq G_q^{(k+1)}$ because strict transforms keep the inclusion relation. Moreover, since $G'_1, \dots, G'_r$ are the minimal elements in $\mathcal{G}^{(k)}$ which contain $Y_k \mathcal{T}$, the subvariety $Y_{k+1} \mathcal{T}$ is the intersection $\cap_{i=1}^r \tilde{G}'_i$. Then

$$\tilde{A} = \bigcap_{i=1}^m \tilde{G}'_i, \quad \tilde{B} = \bigcap_{i=m+1}^r \tilde{G}'_i, \quad (A \cap B)^\sim = \tilde{A} \cap \tilde{B} = \bigcap_{i=1}^m \tilde{G}'_i$$

by [Li06] Lemma 2.9. Thus $Y_{k+1} \mathcal{T} = (Y_k \mathcal{T})^\sim$. By the definition of arrangement we know that $Y_k \mathcal{T}$ and $G_{k+1}^{(k)}$ intersect cleanly, so $Y_{k+1} \mathcal{T}$ is the blow-up of $Y_k \mathcal{T}$ along the center $G_{k+1}^{(k)} \cap Y_k \mathcal{T}$. The exceptional divisor is the preimage of the center, hence is $G_{k+1}^{(k+1)} \cap Y_{k+1} \mathcal{T}$. Since $G_{k+1}^{(k+1)}$ and $Y_{k+1} \mathcal{T}$ intersect cleanly and since the divisor $G_{k+1}^{(k+1)}$ does not contain $Y_{k+1} \mathcal{T}$, we can see that the intersection $G_{k+1}^{(k)} \cap Y_k \mathcal{T}$ is actually transversal.

The codimension of the center $Y_k \mathcal{T} \cap G_{k+1}^{(k)}$ in $Y_k \mathcal{T}$ equals

$$\text{codim}(A \cap B \cap G_{k+1}^{(k)}, A \cap B) = \text{codim}(G_{k+1}^{(k)} \cap B, A \cap B) = \text{codim}(G_{k+1}^{(k)}, A),$$

where the second equality is because of the transversality of the intersection $G_{k+1}^{(k)} \cap B$. If no elements in $\mathcal{T}$ contain G_{k+1} , then $A = Y$ and

$$\text{codim}(G_{k+1}^{(k)}, A) = \dim Y - \dim G_{k+1};$$

otherwise

$$\begin{aligned} \text{codim}(G_{k+1}^{(k)}, A) &= \text{codim}(G_{k+1}^{(k)}, \bigcap_{i=1}^m G'_i) = \text{codim}(G_{k+1}, \bigcap_{G_{k+1} \subsetneq G \in \mathcal{T}} G) \\ &= \dim \bigcap_{G_{k+1} \subsetneq G \in \mathcal{T}} G - \dim G_{k+1}. \end{aligned}$$

Thus the proof is complete. $\square$

3. THE MOTIVE OF WONDERFUL COMPACTIFICATIONS

Notations:

- Let Y be a nonsingular quasi-projective variety with an arrangement of subvarieties $\mathcal{S}$. Let $\mathcal{G}$ be a building set with respect to $\mathcal{S}$. Let $Y_{\mathcal{G}}$ be the wonderful compactification. Let $\mathcal{T}$ be a $\mathcal{G}$ -nest.
- For $T \in \mathcal{G}$, define D_T to be the divisor $T^{(N)}$ in $Y_{\mathcal{G}}$. When no confusion arise, we use the same notation D_T for its restriction to a subvariety of $Y_{\mathcal{G}}$.
- Denote $j_{\mathcal{T}} : Y_{\mathcal{G}} \mathcal{T} \rightarrow Y_{\mathcal{G}}$ to be the natural imbedding. Denote $g_{\mathcal{T}} : Y_{\mathcal{G}} \mathcal{T} \rightarrow Y_0 \mathcal{T}$ to be the restriction of the natural morphism $Y_{\mathcal{G}} \rightarrow Y$.

- Suppose $j : B \rightarrow C$ and $g : B \rightarrow D$ are two morphisms of varieties. Denote by $(j, g) : B \rightarrow C \times D$ the composition of the diagonal map Δ with $f \times g$:

$$(j, g) : B \xrightarrow{\Delta} B \times B \xrightarrow{f \times g} C \times D.$$

- Given $a \in A(P)$, denote by $\{a\}_i$ the image of the projection $A(P) \rightarrow A^i(P)$ of the Chow ring to its degree i direct summand, i.e., taking the codimension i part of a .
- We set $\bigcap_{G \subsetneq T \in \mathcal{T}} T = Y$ if no T satisfies $G \subsetneq T \in \mathcal{T}$. Define

$$r_G := \dim\left(\bigcap_{G \subsetneq T \in \mathcal{T}} T\right) - \dim G.$$

Define

$$N_G := N_G\left(\bigcap_{G \subsetneq T \in \mathcal{T}} T\right)|_{Y_0\mathcal{T}},$$

the restriction to $Y_0\mathcal{T}$ of the normal bundle of G in the ambient space $(\bigcap_{G \subsetneq T \in \mathcal{T}} T)$. Define

$$M_{\mathcal{T}} := \{\underline{\mu} = \{\mu_G\}_{G \in \mathcal{G}} : 1 \leq \mu_G \leq r_G - 1, \mu_G \in \mathbb{Z}\}$$

and define $\|\underline{\mu}\| := \sum_{G \in \mathcal{G}} \mu_G$ for $\underline{\mu} \in M_{\mathcal{T}}$.

Theorem 3.1. *We have the Chow group decomposition*

$$A^*Y_{\mathcal{G}} = A^*Y \oplus \bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}} A^{*- \|\underline{\mu}\|}(Y_0\mathcal{T})$$

where $\mathcal{T}$ runs through all $\mathcal{G}$ -nests.

Moreover, when Y is complete, we have the Chow motive decomposition

$$h(Y_{\mathcal{G}}) = h(Y) \oplus \bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}} h(Y_0\mathcal{T})(\|\underline{\mu}\|)$$

where $\mathcal{T}$ runs through all $\mathcal{G}$ -nests.

Theorem 3.2. *The correspondence that gives each of the above direct summand can be explicitly expressed as follows,*

$$\alpha : h(Y_{\mathcal{G}}) \rightarrow h(Y_0\mathcal{T})(\|\underline{\mu}\|)$$

$$\alpha = (j_{\mathcal{T}}, g_{\mathcal{T}})_* \prod_{G \in \mathcal{T}} \left\{ c\left(g_{\mathcal{T}}^*(N_G) \otimes \mathcal{O}\left(-\sum_{(\star)} D_{G'}\right)\right) \frac{1}{1 + D_G} \right\}_{r_G - 1 - \mu_G},$$

here c is total Chern class, the subscript $r_G - 1 - \mu_G$ means the codimension $r_G - 1 - \mu_G$ part, and condition $(\star)$ is: $G' \subsetneq G$ and $\mathcal{T} \cup \{G'\}$ is a $\mathcal{G}$ -nest.

The inverse correspondence is

$$\beta : h(Y_0\mathcal{T})(\|\underline{\mu}\|) \rightarrow h(Y_{\mathcal{G}})$$

$$\beta = (g_{\mathcal{T}}, j_{\mathcal{T}})_* \prod_{G \in \mathcal{T}} (-D_G)^{\mu_G - 1}.$$

3.1. Proof of the Theorem 3.1.

Lemma 3.3. *Given a $\mathcal{G}$ -nest $\mathcal{T} \subseteq \{G_{k+2}, \dots, G_N\}$. Suppose $\mathcal{T}' := \mathcal{T} \cup \{G_{k+1}\}$ is also a $\mathcal{G}$ -nest. Define $r = r_{k, \mathcal{T}}$ to be*

$$\begin{cases} \dim \cap_{G_{k+1} \subsetneq G \in \mathcal{T}} G - \dim G_{k+1}, & \text{if } \{G : G_{k+1} \subsetneq G \in \mathcal{T}\} \neq \emptyset; \\ \dim Y - \dim G_{k+1}, & \text{otherwise.} \end{cases}$$

Then the following Chow group decomposition holds:

$$A^*(Y_{k+1}\mathcal{T}) = A^*(Y_k\mathcal{T}) \oplus \bigoplus_{t=1}^{r-1} A^{*-t}(Y_k\mathcal{T}').$$

When Y is complete, we also have the motivic decomposition

$$h(Y_{k+1}\mathcal{T}) = h(Y_k\mathcal{T}) \oplus \bigoplus_{t=1}^{r-1} h(Y_k\mathcal{T}')(t).$$

Proof. Apply the well known blow-up formula for the Chow group and for the Chow motive (Theorem A.2) to Proposition 2.7 immediately gives the conclusion. $\square$

Iteratively applying the above lemma gives the proof of Theorem 3.1:

Proof of Theorem 3.1. Define

$$M_{\mathcal{T}}^{(k)} = \{\underline{\mu} = \{\mu_G\}_{G \in \mathcal{G}} : 1 \leq \mu_G \leq \dim(\bigcap_T T^{(k)}) - \dim G^{(k)} - 1, \mu_G \in \mathbb{Z}\}$$

where T runs through the subvarieties in $\mathcal{T}$ such that $G^{(k)} \subsetneq T^{(k)}$. Define $\|\underline{\mu}\| := \sum_{G \in \mathcal{G}} \mu_G$ for $\underline{\mu} \in M_{\mathcal{T}}^{(k)}$.

We prove the following statement using a downward induction on k :

$$(2) \quad A^*Y_{\mathcal{G}} = A^*Y_k \oplus \bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}^{(k)}} A^{*- \|\underline{\mu}\|}(Y_k\mathcal{T}).$$

where $\mathcal{T}$ runs through all $\mathcal{G}$ -nest such that $\mathcal{T} \subseteq \{G_{k+1}, G_{k+2}, \dots, G_N\}$.

The assertion for $k = N$ is trivial because all $G^{(N)}$ are divisors in $Y_{\mathcal{G}}$ hence of codimension 1 and $M_{\mathcal{T}}^{(k)} = \emptyset$.

Assume (2) has been proved for $k+1$, i.e.,

$$A^*Y_{\mathcal{G}} = A^*Y_{k+1} \oplus \bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}^{(k+1)}} A^{*- \|\underline{\mu}\|}(Y_{k+1}\mathcal{T})$$

where $\mathcal{T}$ runs through all $\mathcal{G}$ -nest such that $\mathcal{T} \subseteq \{G_{k+2}, G_{k+3}, \dots, G_N\}$. Apply Lemma 3.3, we have

$$\begin{aligned}
 (3) \quad A^*Y_{\mathcal{G}} = & A^*Y_k \bigoplus \left(\bigoplus_{t=1}^{\text{codim}(G_{k+1}, Y) - 1} A^{*-t}(G_{k+1}^{(k)}) \right) \\
 & \bigoplus \left(\bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}^{(k+1)}} A^{*-||\underline{\mu}||}(Y_k(\mathcal{T})) \right) \\
 & \bigoplus \left(\bigoplus_{\mathcal{T}} \bigoplus_{\underline{\mu} \in M_{\mathcal{T}}^{(k+1)}} \bigoplus_{t=1}^{r_{k+1, \mathcal{T}} - 1} A^{*-||\underline{\mu}|| - t}(Y_k(\{G_{k+1}\} \cup \mathcal{T})) \right).
 \end{aligned}$$

This immediately gives the Chow group decomposition (2) for k . Indeed, any $\mathcal{G}$ -nest contained in $\{G_{k+1}, G_{k+2}, \dots, G_N\}$ must be one of the three: $\{G_{k+1}\}$, a $\mathcal{G}$ -nest $\mathcal{T}$ contained in $\{G_{k+2}, G_{k+3}, \dots, G_N\}$, or $\{G_{k+1}\} \cup \mathcal{T}$. They correspond to the second, third and last summands in (3) respectively. (Notice that $Y_k(\{G_{k+1}\} \cup \mathcal{T}) = \emptyset$ if $\{G_{k+1}\} \cup \mathcal{T}$ is not a $\mathcal{G}$ -nest by Proposition 2.7.)

Therefore, the Chow group decomposition (2) holds for all k , in particular the case $k = 0$ gives the desired Chow group decomposition. For the proof of the Chow motive decomposition, we can either repeat the above proof almost word by word or, as the referee pointed out, notice that the Chow motive decomposition follows from the result on the Chow groups and Manin's identity principle. $\square$

3.2. Proof of Theorem 3.2. First, we introduce some notations. For a given $\mathcal{G}$ -nest $\mathcal{T}$,

- Define $\mathcal{T}_k := \mathcal{T} \cap \{G_{k+1}, G_{k+2}, \dots, G_N\}$ for $0 \leq k \leq N$. Then we have a chain of $\mathcal{G}$ -nests $\mathcal{T}_0 \supseteq \mathcal{T}_1 \supseteq \dots \supseteq \mathcal{T}_N$, where $\mathcal{T}_0 = \mathcal{T}$ and $\mathcal{T}_N = \emptyset$.
- For $\underline{\mu} \in \mathcal{M}_{\mathcal{T}}$ and $1 \leq i \leq N$, define

$$\mu_i := \begin{cases} \mu_{G_i}, & \text{if } G_i \in \mathcal{T}; \\ 0, & \text{otherwise.} \end{cases}$$

- j_{kl} and g_{kl} ($N \geq k > l \geq 0$) are the natural morphisms as in the following diagram

$$(4) \quad \begin{array}{c} \begin{array}{ccccccc} & & & & j_{\mathcal{T}} & & \\ & & & & \curvearrowright & & \\ Y_N \mathcal{T}_0 & \xrightarrow{j_{N0}} & Y_N \mathcal{T}_1 & \xrightarrow{j_{N1}} & \cdots & \xrightarrow{j_{N,N-1}} & Y_N \mathcal{T}_N \\ \downarrow g_{N0} & & \downarrow g_{N1} & & & & \downarrow g_{N,N-1} \\ Y_{N-1} \mathcal{T}_0 & \xrightarrow{j_{N-1,0}} & Y_{N-1} \mathcal{T}_1 & \xrightarrow{j_{N-1,1}} & \cdots & \xrightarrow{j_{N-1,N-1}} & Y_{N-1} \mathcal{T}_{N-1} \\ \downarrow g_{N-1,0} & & \downarrow g_{N-1,1} & & & & \downarrow g_{N-1,N-1} \\ \vdots & & \vdots & & \cdots & & \vdots \\ Y_1 \mathcal{T}_0 & \xrightarrow{j_{10}} & Y_1 \mathcal{T}_1 & & & & \\ \downarrow g_{10} & & \downarrow g_{11} & & & & \\ Y_0 \mathcal{T}_0 & & Y_0 \mathcal{T}_1 & & & & \end{array} \end{array}$$

(Note: The diagram shows a grid of squares. Horizontal arrows are j_{kl} and vertical arrows are g_{kl} . Dotted arrows α_k and β_k connect $Y_k \mathcal{T}_0$ to $Y_k \mathcal{T}_k$ and $Y_{k-1} \mathcal{T}_{k-1}$ respectively. A large curved arrow $j_{\mathcal{T}}$ connects $Y_N \mathcal{T}_0$ to $Y_N \mathcal{T}_N$. A large curved arrow $g_{\mathcal{T}}$ connects $Y_N \mathcal{T}_0$ to $Y_0 \mathcal{T}_0$.)

Lemma 3.4. Denote by $g : Y_k \rightarrow Y_{k-1}$ the natural morphism. Then for $l \leq k-1$, we have

$$g^{-1}(G_l^{(k-1)}) = G_l^{(k)}.$$

Proof. First, we claim that $G_l^{(k-1)} \not\supseteq G_k^{(k-1)}$. Otherwise $G_l \supseteq G_k$ since they are the respective images of $G_l^{(k-1)}$ and $G_k^{(k-1)}$ under $Y_k \rightarrow Y_0$. But then by the assumption that the order of $\{G_i\}$ is compatible with inclusion relations, we obtain a contradiction $l \geq k$.

Next, it is easy to see that $G_l^{(k-1)} \not\subseteq G_k^{(k-1)}$ since $G_l^{(k-1)}$ is a divisor. Now we know that the two nonsingular subvarieties $G_l^{(k-1)}$ and $G_k^{(k-1)}$ intersect cleanly and neither one contains the other, therefore they must intersect transversally. Then it is standard to show by local coordinates calculation that the following isomorphism between ideal sheaves holds:

$$g^{-1}\mathcal{I}(G_l^{(k-1)}) \cdot \mathcal{O}_{Y_k} \cong \mathcal{I}(G_l^{(k)}).$$

The desired conclusion follows from this. $\square$

Lemma 3.5. In Diagram (4), all squares are fiber squares. Moreover, for any $N \geq k > l \geq 0$, we have

- (i) j_{kl} is injective;
- (ii) If $G_k \in \mathcal{T}$, then g_{kl} is the projection of a projective bundle with fiber isomorphic to a projective space of dimension $r_{k,\mathcal{T}} - 1$;
- (iii) If $G_k \notin \mathcal{T}$ but $\{G_k\} \cup \mathcal{T}_l$ is a $\mathcal{G}$ -nest, then g_{kl} is the blow-up of $Y_{k-1}\mathcal{T}_l$ along the center $G_k^{(k-1)} \cap Y_{k-1}\mathcal{T}_l$;
- (iv) If $\{G_k\} \cup \mathcal{T}_l$ is not a $\mathcal{G}$ -nest, then g_{kl} is an isomorphism.

Proof. It is obvious that j_{kl} is injective.

Now we show that g_{kl} is the projection of a projective bundle if $G_k \in \mathcal{T}$. By Proposition 2.7, the variety $Y_k \mathcal{T}_k$ is the blow-up of $Y_{k-1} \mathcal{T}_k$ along the center $Y_{k-1} \mathcal{T}_{k-1}$, and the exceptional divisor is $Y_k \mathcal{T}_{k-1}$ (note that $Y_{k-1} \mathcal{T}_k \cap G_k^{(k-1)} = Y_{k-1} \mathcal{T}_{k-1}$ and $Y_k \mathcal{T}_k \cap G_k^{(k)} = Y_k \mathcal{T}_{k-1}$). Therefore $g_{k,k-1} : Y_k \mathcal{T}_{k-1} \rightarrow Y_{k-1} \mathcal{T}_{k-1}$ is a projective bundle, and the dimension of a fibre is $r_{k,\mathcal{T}} - 1$. Next we show that for any $l \leq k-1$, g_{kl} is the restriction of $g_{k,k-1}$ to a smaller base $Y_{k-1} \mathcal{T}_l$, which will then show that g_{kl} is also a projective bundle with fiber of the same dimension $r_{k,\mathcal{T}} - 1$. Fix k and use downward induction on l . By inductive assumption, $g_{k,l+1}$ is a restriction of $g_{k,k-1}$. Since

$$g_{k,l+1}^{-1}(G_{l+1}^{(k-1)} \cap Y_{k-1} \mathcal{T}_l) = G_{l+1}^{(k)} \cap Y_k \mathcal{T}_l$$

by Lemma 3.4, the restriction of the projective bundle $g_{k,l+1}$ to a smaller base space $Y_{k-1} \mathcal{T}_l = Y_{k-1} \mathcal{T}_{l+1} \cap G_{l+1}^{(k-1)}$ is exactly g_{kl} .

Next, we show g_{kl} is birational if $G_k \notin \mathcal{T}$. This is again implied by Proposition 2.7. Notice that $G_k^{(k-1)}$ is minimal in

$$\mathcal{T}' := \{G_k^{(k-1)}\} \cup \{G^{(k-1)}\}_{G \in \mathcal{T}_l}.$$

If $\mathcal{T}'$ is a $\mathcal{G}^{(k-1)}$ -nest, then $g_{kl} : Y_k \mathcal{T}_l \rightarrow Y_{k-1} \mathcal{T}_l$ is a blow-up along the center $G_k^{(k-1)} \cap Y_{k-1} \mathcal{T}_l$; otherwise, g_{kl} is an isomorphism. In both cases, g_{kl} is birational.

Finally, all squares in Diagram (4) are fiber squares since $\forall l \leq k-2$, g_{kl} is a restriction of $g_{k,l+1}$. The proof is complete. $\square$

The following lemma computes the composition of correspondences in certain diagrams. The author thanks the referee to suggest a proof much simpler than the original proof given by the author.

Lemma 3.6. *Let W, U, V, X, Y, Z be nonsingular quasi-projective varieties. Suppose the square in the following diagram is a fiber square.*

$$\begin{array}{ccccc} W & \xrightarrow{j_3} & U & \xrightarrow{j_2} & X \\ g_3 \downarrow & \square & g_2 \downarrow & \nearrow \alpha_2 & \nearrow \beta_2 \\ V & \xrightarrow{j_1} & Y & & \\ g_1 \downarrow & \nearrow \alpha_1 & \nearrow \beta_1 & & \\ Z & & & & \end{array}$$

and suppose that $\dim W - \dim V = \dim U - \dim Y$ and that $j_k, g_k (1 \leq k \leq 3)$ are proper. Take $\gamma_1, \gamma'_1 \in A(V)$, $\gamma_2, \gamma'_2 \in A(U)$ and define correspondences

$$\alpha_k = (j_k, g_k)_* \gamma_k, \quad \beta_k = (g_k, j_k)_* \gamma'_k, \quad \text{for } k = 1, 2.$$

Then we have

$$(5) \quad \alpha_1 \alpha_2 = (j_2 j_3, g_1 g_3)_* (j_3^* \gamma_2 \cdot g_3^* \gamma_1),$$

$$(6) \quad \beta_2 \beta_1 = (g_1 g_3, j_2 j_3)_* (g_3^* \gamma'_1 \cdot j_3^* \gamma'_2).$$

Proof. By abuse of notation, for $\gamma \in A(V)$ we use the same γ to denote the correspondence $(\Delta_V)_*(\gamma) \in A(V \times V)$ where $\Delta_V : V \rightarrow V \times V$ is the diagonal embedding. For a map $j : U \rightarrow X$, we denote by j_* the correspondence Γ_j (i.e. the graph of j) and by j^* the correspondence Γ'_j (i.e. the transpose of Γ_j).

First observe that $\alpha_k = g_{k*} \circ \gamma \circ j_k^*$ for $k = 1, 2$. Indeed, by properties of correspondences (cf. [Fu98] Prop 16.1.1(c)), we have $\Gamma_j \circ \gamma = (1_U \times j)_* \gamma$, $\gamma \circ \Gamma'_j = (g \times 1_U)_* \gamma$, so

$$g_{k*} \circ \gamma \circ j_k^* = \Gamma_{g_k} \circ \gamma_k \circ \Gamma_{j_k}^* = (g_k \times j_k)_* \gamma_k = (g_k, j_k)_* \gamma_k = \alpha_k \quad \text{for } k = 1, 2.$$

With the above observation, (5) is equivalent to

$$g_{1*} \gamma_1 j_1^* g_{2*} \gamma_2 j_2^* = (g_1 g_3)_* (j_3^* \gamma_2 \cdot g_3^* \gamma_1) (j_2 j_3)^*.$$

So it suffices to prove

$$(7) \quad \gamma_1 j_1^* g_{2*} \gamma_2 = g_{3*} (j_3^* \gamma_2 \cdot g_3^* \gamma_1) j_3^*.$$

For any $u \in A(U)$, we have

$$\gamma_1 j_1^* g_{2*} \gamma_2(u) = \gamma_1 g_{3*} j_3^* \gamma_2(u) = g_{3*} (g_3^* \gamma_1 \cdot j_3^* (\gamma_2 u)) = g_{3*} (g_3^* \gamma_1 \cdot j_3^* \gamma_2) j_3^*(u)$$

where the first “=” is because of $\dim W - \dim V = \dim U - \dim Y$, the second “=” is because of the projection formula. Then we apply Manin’s Identity Principle to obtain (7), hence (5). The identity (6) can be obtained by transposing (5). $\square$

Now we state a simple lemma and omit the proof.

Lemma 3.7. *If A, B_i, C_{ij} are motives such that*

- (i) $\bigoplus_i \alpha_i : A \cong \bigoplus_i B_i$ *is an isomorphism with inverse $\sum_i \beta_i$, and*
- (ii) $\bigoplus_j \alpha_{ij} : B_i \cong \bigoplus_j C_{ij}$ *is an isomorphism with inverse $\sum_j \beta_{ij}$,*

then the correspondence $\bigoplus_{i,j} \alpha_{ij} \circ \alpha_i$ gives an isomorphism $A \cong \bigoplus_{i,j} C_{ij}$ with inverse $\sum_{i,j} \beta_i \circ \beta_{ij}$.

For $G_k \in \mathcal{T}$, define $h_k \in A^1(Y_k \mathcal{T}_{k-1})$ to be first Chern class of the invertible sheaf $\mathcal{O}(1)$ of the projective bundle $g_{k,k-1}$. Define

$$\alpha_k = \begin{cases} (j_{k,k-1}, g_{k,k-1})_* 1, & \text{if } G_k \notin \mathcal{T}; \\ (j_{k,k-1}, g_{k,k-1})_* \left(\{g_{k,k-1}^* c(N_k) \frac{1}{1-h_k}\}_{r_{k-1}-\mu_k} \right), & \text{if } G_k \in \mathcal{T}, \end{cases}$$

where $N_k := N_{Y_{k-1} \mathcal{T}_{k-1}} Y_{k-1} \mathcal{T}_k$. Define

$$\beta_k = \begin{cases} (g_{k,k-1}, j_{k,k-1})_* 1, & \text{if } G_k \notin \mathcal{T}; \\ (g_{k,k-1}, j_{k,k-1})_* h_k^{\mu_k-1}, & \text{if } G_k \in \mathcal{T}. \end{cases}$$

Thanks to the blow-up formula of motives (Theorem A.2), the correspondence

$$a_k : h(Y_k \mathcal{T}_k) \left(\sum_{i=k+1}^N \mu_i \right) \rightarrow h(Y_{k-1} \mathcal{T}_{k-1}) \left(\sum_{i=k}^N \mu_i \right)$$

expresses $h(Y_{k-1}\mathcal{T}_{k-1})(\sum_k^N \mu_i)$ as a direct summand of $h(Y_k\mathcal{T}_k)(\sum_{k+1}^N \mu_i)$ with right inverse β_k .

By Lemma 3.7, the correspondence

$$\alpha_{\mathcal{T}, \underline{\mu}} : h(Y_{\mathcal{G}}) \rightarrow h(Y_0\mathcal{T})(\|\underline{\mu}\|)$$

that gives the direct summand $h(Y_0\mathcal{T})(\|\underline{\mu}\|)$ in Theorem 3.1 can be expressed as the composition $\alpha_1 \circ \alpha_2 \circ \cdots \circ \alpha_N$, with right inverse $\beta_N \circ \cdots \circ \beta_1$. Therefore we have

Proposition 3.8. *Denote by $f_k : Y_N\mathcal{T}_0 \rightarrow Y_k\mathcal{T}_{k-1}$ the natural map in Diagram (4). (i.e. $g_{k+1,k-1} \cdots \circ g_{N,k-1} \circ j_{N,k-2} \circ \cdots \circ j_{N0}$.) Then*

$$\begin{aligned} \alpha_1 \circ \cdots \circ \alpha_N &= (j_{\mathcal{T}}, g_{\mathcal{T}})_* \prod_{G_k \in \mathcal{T}} \left\{ f_k^* g_{k,k-1}^* c(N_k) \frac{1}{1 - f_k^* h_k} \right\}_{r_k-1-\mu_k}, \\ \beta_N \circ \cdots \circ \beta_1 &= (g_{\mathcal{T}}, j_{\mathcal{T}})_* \prod_{G_k \in \mathcal{T}} f_k^* h_k^{\mu_k-1}. \end{aligned}$$

Proof. Combine Lemma 3.5 and Lemma 3.6 with the above discussion. $\square$

The following two standard facts about normal bundles of subvarieties are used in the proof of Theorem 3.2.

Fact 3.9. *Let Z be a nonsingular variety. Let Y, W be nonsingular proper subvarieties of Z and assume Y intersects transversally with W . Let $\pi : \tilde{Z} \rightarrow Z$ be the blow-up of Z along W and let $\tilde{Y}$ be the strict transform of Y . Then*

$$N_{\tilde{Y}} \tilde{Z} \simeq \pi^* N_Y Z.$$

Fact 3.10. *Let $W \subsetneq Y \subsetneq Z$ be nonsingular varieties and $\pi : \tilde{Z} \rightarrow Z$ be the blow-up of Z along W . Denote by $\tilde{Y}$ the strict transform of Y , and denote by E the exceptional divisor on $\tilde{Y}$. Then*

$$N_{\tilde{Y}} \tilde{Z} \simeq \pi^* N_Y Z \otimes \mathcal{O}(-E).$$

Proof of the above two facts. Prove by local coordinates. Or see [Fu98]. $\square$

Proof of Theorem 3.2. To conclude Theorem 3.2 from Proposition 3.8, we prove in three steps.

Step 1: Show $f_k^* h_k = -D_{G_k}|_{Y_N\mathcal{T}_0}$.

Recall that for $G_k \in \mathcal{T}$, h_k is first Chern class of the invertible sheaf $\mathcal{O}(1)$ of the projective bundle $g_{k,k-1}$.

Consider the following diagram (not necessary a fiber square) where π and j are the natural morphisms:

$$\begin{array}{ccc} Y_N\mathcal{T}_0 & \xrightarrow{j_{\mathcal{T}}} & Y_N \\ \downarrow f_k & & \downarrow \pi \\ Y_k\mathcal{T}_{k-1} & \xrightarrow{j} & Y_k. \end{array}$$

By Proposition 2.7, $Y_k \mathcal{T}_{k-1}$ is the exceptional divisor of the blow-up $g_{k,k-1} : Y_k \mathcal{T}_{k-1} \rightarrow Y_{k-1} \mathcal{T}_{k-1}$, so $h_k = -j_{k,k-1}^*[Y_k \mathcal{T}_{k-1}]$. Since $Y_k \mathcal{T}_{k-1}$ is the transversal intersection $Y_k \mathcal{T}_k \cap G_k^{(k)}$, $h_k = -j^*[G_k^{(k)}]$. Then

$$f_k^* h_k = -f_k^* j^*[G_k^{(k)}] = -j_T^* \pi^*[G_k^{(k)}] = -j_T^* D_{G_k} = -D_{G_k}|_{Y_N \mathcal{T}_0}.$$

where the third equality can be proved by successively applying Lemma 3.4.

Step 2: Let $0 \leq s < k \leq N$. Denote $g_{sk} : Y_s \mathcal{T}_k \rightarrow Y_{s-1} \mathcal{T}_k$ to be the natural map induced from $Y_s \rightarrow Y_{s-1}$. We claim the following:

If $G_k \in \mathcal{T}$ (hence $\mathcal{T}_{k-1} = \mathcal{T}_k \cup \{G_k\}$), then the normal bundle $N_{Y_s \mathcal{T}_{k-1}} Y_s \mathcal{T}_k$ is isomorphic to

$$\begin{cases} g_{s,k-1}^*(N_{Y_{s-1} \mathcal{T}_{k-1}} Y_{s-1} \mathcal{T}_k) \otimes (-[G_s^{(s)}]|_{Y_s \mathcal{T}_{k-1}}), & \text{if } (**) \text{ holds;} \\ g_{s,k-1}^*(N_{Y_{s-1} \mathcal{T}_{k-1}} Y_{s-1} \mathcal{T}_k), & \text{otherwise.} \end{cases}$$

where condition $(**)$ is: $G_s \subsetneq G_k$ and $\mathcal{T}_k \cup \{G_s\}$ is a $\mathcal{G}$ -nest.

For the proof, we discuss three cases.

Case (i): condition $(**)$ holds. It is a direct conclusion of Fact 3.10. Indeed, to apply Fact 3.10 we need to show that

$$Y_{s-1} \mathcal{T}_k \cap G_s^{(s-1)} \subsetneq Y_{s-1} \mathcal{T}_k \cap G_k^{(s-1)} \subsetneq Y_{s-1} \mathcal{T}_k.$$

The second inequality is obvious. The first inclusion is strict because of the following reason. $G_s^{(s-1)}$ is a $\mathcal{G}^{(s-1)}$ -factor of $Y_{s-1} \mathcal{T}_k \cap G_s^{(s-1)}$, therefore $G_k^{(s-1)}$ is not a $\mathcal{G}^{(s-1)}$ -factor because it strictly contains $G_s^{(s-1)}$. On the other hand, $G_k^{(s-1)}$ is a $\mathcal{G}^{(s-1)}$ -factor of $Y_{s-1} \mathcal{T}_k \cap G_k^{(s-1)}$. So the first inclusion is strict.

Case (ii): $\mathcal{T}_k \cup \{G_s\}$ is not $\mathcal{G}$ -nested. In this case, $G_s^{(s-1)} \cap Y_{s-1} \mathcal{T}_k = \emptyset$ by Proposition 2.7. Hence no twisting is needed for the normal bundle.

Case (iii): $\mathcal{T}_k \cup \{G_s\}$ is $\mathcal{G}$ -nested but G_s is not strictly contained in G_k . If $\mathcal{T}_{k-1} \cup \{G_s\}$ is not a $\mathcal{G}$ -nest, then $G_s^{(s-1)} \cap Y_{s-1} \mathcal{T}_{k-1} = \emptyset$ by Proposition 2.7. Hence blowing up along $G_s^{(s-1)}$ will not affect the normal bundle of $Y_{s-1} \mathcal{T}_{k-1}$, so no twisting is needed. Otherwise, assume $\mathcal{T}_{k-1} \cup \{G_s\}$ is a $\mathcal{G}$ -nest. Both G_s and G_k are minimal in the $\mathcal{G}$ -nest $\mathcal{T}_{k-1} \cup \{G_s\}$. Then $G_s^{(s-1)}$ and $G_k^{(s-1)}$ are minimal in a nest and neither one contains the other, therefore they intersect transversally by the definition of nest. Thus, $Y_{s-1} \mathcal{T}_k \cap G_k^{(s-1)}$ and $Y_{s-1} \mathcal{T}_k \cap G_s^{(s-1)}$, regarded as subvarieties of the ambient space $Y_{s-1} \mathcal{T}_k$, intersect transversally. Therefore Fact 3.9 applies, and no twisting is needed for the normal bundle.

Step 3: Apply the result of Step 2 successively for $s = 1, 2, \dots, k-1$. The normal bundle $N_{Y_{k-1} \mathcal{T}_{k-1}} Y_{k-1} \mathcal{T}_k$ is isomorphic to

$$\left(g_{k-1,k-1}^* \cdots g_{1,k-1}^* (N_{Y_0 \mathcal{T}_{k-1}} Y_0 \mathcal{T}_k) \right) \otimes \left(- \sum_{(**)} [G_s^{(k-1)}]|_{Y_{k-1} \mathcal{T}_{k-1}} \right)$$

where the sum is over all s that satisfying condition (**). (Here we have used Lemma 3.4.) Therefore

$$\begin{aligned} & f_k^* g_{k,k-1}^* c(N_{Y_{k-1}\mathcal{T}_{k-1}} Y_{k-1}\mathcal{T}_k) \\ &= c\left(g_{\mathcal{T}}^*(N_{Y_0\mathcal{T}_{k-1}} Y_0\mathcal{T}_k|_{Y_0\mathcal{T}}) \otimes \mathcal{O}\left(-\sum_{(**)} [D_{G_s}]|_{Y_N\mathcal{T}_{k-1}}\right)\right). \end{aligned}$$

Notice that

$$(N_{Y_0\mathcal{T}_{k-1}} Y_0\mathcal{T}_k)|_{Y_0\mathcal{T}} = N_{G_k}\left(\bigcap_{G_k \subsetneq G \in \mathcal{T}} G\right)|_{Y_0\mathcal{T}}$$

which is denoted by N_{G_k} by our notation. (The proof is as follows: Suppose $T_1, \dots, T_m, T_{m+1}, \dots, T_r$ are the minimal elements of the nest $\mathcal{T}_k$, where the first m elements contain G_k . Then the minimal element of the nest $\mathcal{T}_{k-1}$ are $G_k, T_{m+1}, \dots, T_r$. By the definition of nest, $Y_0\mathcal{T}_k$ is the transversal intersection $T_1 \cap \dots \cap T_m \cap T_{m+1} \cap \dots \cap T_r$, and $Y_0\mathcal{T}_{k-1}$ is the transversal intersection $G_k \cap T_{m+1} \cap \dots \cap T_r$. Therefore

$$N_{Y_0\mathcal{T}_{k-1}} Y_0\mathcal{T}_k = N_{G_k}(T_1 \cap \dots \cap T_m)|_{Y_0\mathcal{T}_{k-1}}.$$

Since $T_1 \cap \dots \cap T_m = \bigcap_{G_k \subsetneq G \in \mathcal{T}} G$, the conclusion follows immediately.)

Now put everything into Corollary 3.8, we have

$$\begin{aligned} & \alpha_1 \circ \dots \circ \alpha_N \\ &= (j_{\mathcal{T}}, g_{\mathcal{T}})_* \prod_{G_k \in \mathcal{T}} \left\{ c(g_{\mathcal{T}}^*(N_{G_k}) \otimes \mathcal{O}\left(-\sum_{(*)'} [D_{G_s}]|_{Y_N\mathcal{T}}\right) \frac{1}{1 + D_{G_k}|_{Y_N\mathcal{T}}}) \right\}_{r_k-1-\mu_k}, \\ & \beta_N \circ \dots \circ \beta_1 = (g_{\mathcal{T}}, j_{\mathcal{T}})_* \prod_{G_k \in \mathcal{T}} (-D_{G_k})^{\mu_k-1}|_{Y_N\mathcal{T}}. \end{aligned}$$

Finally, we show that the condition (**) can be replaced by the following condition:

($\star$): $G_s \subsetneq G_k$ and $\mathcal{T} \cup \{G_s\}$ is a $\mathcal{G}$ -nest.

Indeed, ($\star$) is stronger than (**). However, for those G_s satisfying (**) but not ($\star$), the divisor $[D_{G_s}]|_{Y_N\mathcal{T}}$ would be trivial because $D_{G_s} \cap Y_N\mathcal{T} = \emptyset$. Therefore, replacing (**) by ($\star$) will not affect the resulting correspondence.

Hence the proof is complete. $\square$

We write a direct conclusion from Step 3 for later usage:

Corollary 3.11. Denote $\pi : G_{k+1}^{(k)} \rightarrow G_{k+1}$. Then

$$c(N_{G_{k+1}^{(k)}} Y_k) = c\left(\pi^* N_{(G_{k+1})} Y \otimes \sum_{G_{k+1} \supsetneq G \in \mathcal{T}} (-[D_G])|_{G_{k+1}^{(k)}}\right).$$

Proof. Apply Step 3 to the nest $\mathcal{T} = \{G_{k+1}\}$. $\square$

4. FULTON-MACPHERSON CONFIGURATION SPACES

Fix a nonsingular variety X of dimension d . The configuration space of n distinct ordered points on X , denoted by $F(X, n)$, can be naturally identified with an open subvariety of the Cartesian product X^n :

$$F(X, n) := \{(x_1, x_2, \dots, x_n) \in X^n : x_i \neq x_j \text{ for } i \neq j\}.$$

In their celebrated paper [FM94], Fulton and MacPherson have discovered an interesting compactification $X[n]$ of the configuration space $F(X, n)$. The compactification is obtained by replacing the diagonals of X^n by a simple normal crossing divisor. It has many attractive properties, for example the geometry when n points collide, i.e. the degenerate configuration, can be explicitly described using $X[n]$. $X[n]$ is closely related to the well known compactification $\overline{\mathcal{M}}_{0,n}$ of the moduli space of stable rational curves with n marked points. The reader is referred to the beautiful paper [FM94] for the original construction and various applications of the Fulton-MacPherson configuration space.

The Fulton-MacPherson configuration space $X[n]$ can be realized as a wonderful compactification of an arrangement of subvarieties by taking $Y = X^n$, $\mathcal{G}$ the collection of all diagonals of X^n and therefore the induced arrangement is the set of intersections of diagonals which is called polydiagonals (cf. [Li06]).


4.1. Main theorems. First we fix some notations:

(i) Denote $[n] := \{1, 2, \dots, n\}$. We call two subsets $I, J \subseteq [n]$ *overlapped* if $I \cap J$ is a nonempty proper subset of I and J . For a set $\mathcal{S}$ of subsets of $[n]$, we call I is compatible with $\mathcal{S}$ (denote by $I \sim \mathcal{S}$) if I does not overlap any element in $\mathcal{S}$.

A *nest* $\mathcal{S}$ is a set of subsets of $[n]$ such that any two elements $I \neq J \in \mathcal{S}$ are not overlapped, and all singletons $\{1\}, \dots, \{n\}$ are in $\mathcal{S}$. Notice that the nest defined here, unlike the one defined in [FM94], is allowed to contain singletons.

Given a nest $\mathcal{S}$, define $\mathcal{S}^\circ = \mathcal{S} \setminus \{\{1\}, \dots, \{n\}\}$. In the description of nests by forests below, $\mathcal{S}^\circ$ correspond to the forest $\mathcal{S}$ cutting of all leaves.

A nest $\mathcal{S}$ naturally corresponds to forest (i.e. a not necessarily connected tree), each node of which is labeled by an element in $\mathcal{S}$. For example, the following forest corresponds to a nest $\mathcal{S} = \{1, 2, 3, 23, 123\}$.



Denote by $c(\mathcal{S})$ the number of connected components of the forest, i.e., the number of maximal elements of $\mathcal{S}$. Denote by $c_I(\mathcal{S})$ (or c_I if no ambiguity arise) the number of maximal elements of the set $\{J \in \mathcal{S} | J \subsetneq I\}$, i.e. the

number of sons of the node I . In the above example, $c(\mathcal{S}) = 1$, $c_{123} = c_{23} = 2$.

(ii) For a subset $I \subseteq [n]$ consisting of at least two elements, define the diagonal

$$\Delta_I := \{(x_1, \dots, x_n) \in X^n : x_i = x_j \text{ if } i, j \in I\}.$$

It is shown in [FM94] that complement of $F(x, n)$ in the Fulton-MacPherson compactification $X[n]$ is a union of normal crossing nonsingular divisors D_I , indexed by subsets $I \subseteq [n]$ with at least two elements. More precisely, D_I is the dominant transform $\tilde{\Delta}_I$ under the natural morphism $X[n] \rightarrow X^n$.

For every nest $\mathcal{S}$, $X(\mathcal{S}) := \cap_{I \in \mathcal{S}} D_I$ is a nonsingular subvariety of $X[n]$. Define $j_{\mathcal{S}} : X(\mathcal{S}) \hookrightarrow X[n]$ to be the natural inclusion.

Define $\Delta_{\mathcal{S}} := \cap_{I \in \mathcal{S}} \Delta_I$. Define $g_{\mathcal{S}} : X(\mathcal{S}) \rightarrow \Delta_{\mathcal{S}}$ to be the restriction of the morphism $\pi : X[n] \rightarrow X^n$ to the subvariety $X(\mathcal{S})$.

(iii) Let $p_I : X[n] \rightarrow X$ be the composition of $\pi : X[n] \rightarrow X^n$ with the projection $X^n \rightarrow X$ to the i -th factor for an arbitrary $i \in I$. (The choice of $i \in I$ is not essential: indeed, the only place we need p_I is in the formulation of $\alpha_{\mathcal{S}, \mu}$ below, where we need the composition $j_{\mathcal{S}}^* p_I^*$. By the following diagram

$$\begin{array}{ccc} X(\mathcal{S}) & \xrightarrow{j_{\mathcal{S}}} & X[n] \\ \downarrow g_{\mathcal{S}} & & \downarrow p_i \\ \Delta_{\mathcal{S}} & \xrightarrow{q_i} & X \end{array}$$

where $i \in I$, we have $j_{\mathcal{S}}^* p_i^* = g_{\mathcal{S}}^* q_i^*$, but q_i is independent of the choice of $i \in I$ since $\Delta_{\mathcal{S}} \subseteq \Delta_I$, so $j_{\mathcal{S}}^* p_I^*$ is independent of the choice of $i \in I$ for p_I .)

(iv) For a nest $\mathcal{S} \neq \{\{1\}, \dots, \{n\}\}$ (i.e. $\mathcal{S}^\circ \neq \emptyset$), define

$$M_{\mathcal{S}} := \{\underline{\mu} = \{\mu_I\}_{I \in \mathcal{S}^\circ} : 1 \leq \mu_I \leq d(c_I - 1) - 1, \mu_I \in \mathbb{Z}\}.$$

(recall that $d = \dim X$ and $c_I = c_I(\mathcal{S})$ is defined in (i)) and define

$$\|\underline{\mu}\| := \sum_{I \in \mathcal{S}^\circ} \mu_I, \quad \forall \underline{\mu} \in M_{\mathcal{S}}.$$

For $\mathcal{S} = \{\{1\}, \dots, \{n\}\}$, assume $M_{\mathcal{S}} = \{\underline{\mu}\}$ with $\|\underline{\mu}\| = 0$.

We will show in the proof of Theorem 4.1 that $M_{\mathcal{S}}$ is the special case of $M_{\mathcal{T}}$ defined in §3 where Y is X^n , $\mathcal{G}$ is the set of diagonals of X^n and $\mathcal{T}$ is the set of $\mathcal{G}$ -nests.

Define function $\zeta(x) := \sum_{i=0}^d (1+x)^{d-i} c_i(T_X)$.

Define $\alpha_{\mathcal{S}, \underline{\mu}} \in \text{Corr}^{-\|\underline{\mu}\|}(X[n], \Delta_{\mathcal{S}})$, $\beta_{\mathcal{S}, \underline{\mu}} \in \text{Corr}^{\|\underline{\mu}\|}(\Delta_{\mathcal{S}}, X[n])$, $p_{\mathcal{S}, \underline{\mu}} \in \text{Corr}^0(X[n], X[n])$ as follows,

$$\alpha_{\mathcal{S}, \underline{\mu}} = (j_{\mathcal{S}}, g_{\mathcal{S}})_* j_{\mathcal{S}}^* \left(\prod_{I \in \mathcal{S}^\circ} \left\{ -p_I^* \zeta \left(- \sum_{\substack{J \sim \mathcal{S} \\ J \supsetneq I}} D_J \right)^{c_I-1} \frac{1}{1+D_I} \right\}_{d(c_I-1)-1-\mu_I} \right),$$

$$\beta_{\mathcal{S}, \underline{\mu}} = (g_{\mathcal{S}}, j_{\mathcal{S}})_* j_{\mathcal{S}}^* \left(\prod_{I \in \mathcal{S}^\circ} D_I^{\mu_I-1} \right),$$

$$p_{\mathcal{S}, \underline{\mu}} = \beta_{\mathcal{S}, \underline{\mu}} \circ \alpha_{\mathcal{S}, \underline{\mu}}.$$

(In the above definition of $\alpha_{\mathcal{S}, \underline{\mu}}$ and $\beta_{\mathcal{S}, \underline{\mu}}$, the products are set to be $1_{X(\mathcal{S})} \in A^0(X(\mathcal{S}))$ if $\mathcal{S}^\circ = \emptyset$.)

The following are the main theorems on the Chow groups and Chow motives of Fulton-MacPherson configuration spaces.

Theorem 4.1. *Let X be a nonsingular quasi-projective variety. There is an isomorphism of Chow groups:*

$$A^*(X[n]) = \bigoplus_{\mathcal{S}} \bigoplus_{\underline{\mu} \in M_{\mathcal{S}}} A^{*- \|\underline{\mu}\|}(X^{c(\mathcal{S})}).$$

where $\mathcal{S}$ runs through all nests of $[n]$.

Theorem 4.2. *Let X be a nonsingular projective variety. Then there is a canonical isomorphism of Chow motives*

$$\bigoplus_{\mathcal{S}} \bigoplus_{\underline{\mu} \in M_{\mathcal{S}}} \alpha_{\mathcal{S}, \underline{\mu}} : h(X[n]) \cong \bigoplus_{\mathcal{S}} \bigoplus_{\underline{\mu} \in M_{\mathcal{S}}} h(\Delta_{\mathcal{S}})(\|\underline{\mu}\|)$$

with the inverse $\sum_{\mathcal{S}} \sum_{\underline{\mu} \in \mathcal{S}} \beta_{\mathcal{S}, \underline{\mu}}$. Equivalently, we have

$$h(X[n]) \cong \bigoplus_{\mathcal{S}} \bigoplus_{\underline{\mu} \in M_{\mathcal{S}}} h(X^{c(\mathcal{S})})(\|\underline{\mu}\|).$$

Remark: Observe that the two sets of correspondences $\{\alpha_{\mathcal{S}, \underline{\mu}}\}$, $\{\beta_{\mathcal{S}, \underline{\mu}}\}$ are $\mathfrak{S}_n$ -symmetric in the sense that the following diagram commutes for any $\sigma \in \mathfrak{S}_n$,

$$\begin{array}{ccccc} h(X[n]) & \xrightarrow{\alpha_{\mathcal{S}, \underline{\mu}}} & h(\Delta_{\mathcal{S}})(\|\underline{\mu}\|) & \xrightarrow{\beta_{\mathcal{S}, \underline{\mu}}} & h(X[n]) \\ \downarrow \sigma & & \downarrow \sigma & & \downarrow \sigma \\ h(X[n]) & \xrightarrow{\alpha_{\sigma(\mathcal{S}), \underline{\mu}}} & h(\Delta_{\sigma(\mathcal{S})})(\|\underline{\mu}\|) & \xrightarrow{\beta_{\sigma(\mathcal{S}), \underline{\mu}}} & h(X[n]). \end{array}$$

Proof of Theorem 4.1. Apply Theorem 3.1 with the ambient space $Y = X^n$ and the building set

$$\mathcal{G} = \{\Delta_I\}_{I \subseteq [n], |I| \geq 2}$$

First notice that a nest $\mathcal{S}$ of $[n]$ gives a $\mathcal{G}$ -nest $\mathcal{T} = \{\Delta_I\}_{I \in \mathcal{S}^\circ}$. Moreover, the inverse is also true: a $\mathcal{G}$ -nest will give a nest of $[n]$. Indeed, given a partition $\Pi = (I_1, \dots, I_t)$ of $[n]$, a $\mathcal{G}$ -factor of Δ_Π by definition is a minimal element in $\{G \in \mathcal{G} : G \supseteq \Delta_\Pi\}$. So $\{\Delta_{I_1}, \dots, \Delta_{I_t}\}$ are all the $\mathcal{G}$ -factors of Δ_Π . By the definition of $\mathcal{G}$ -nest, $\mathcal{T}$ is induced from a flag of strata

$$\Delta_{\Pi_1} \supseteq \Delta_{\Pi_2} \supseteq \dots \supseteq \Delta_{\Pi_t}.$$

Then

$$\Pi_1 \geq \Pi_2 \geq \dots \geq \Pi_t.$$

(Here $\Pi \geq \Pi'$ means Π is a finer partition than Π' , e.g. $(12, 3, 4) \geq (123, 4)$.) The nest $\mathcal{T}$ is induced by “taking the union of all factors of each Δ_Π ”, which corresponds to “taking all I ’s that appear in any of the partition Π_i ”. Since the partitions are totally ordered, the set of I ’s forms a nest of $[n]$.

Next we prove the range of $\underline{\mu}$ is as stated. Theorem 3.1 asserts that

$$1 \leq \mu_G \leq r_G - 1.$$

Now $G = \Delta_I$ is a diagonal, so by definition

$$\begin{aligned} r_G &:= \dim\left(\bigcap_{G \subsetneq T \in \mathcal{T}} T\right) - \dim G \\ &= \dim\left(\bigcap_{I \supsetneq I' \in \mathcal{S}} \Delta_{I'}\right) - \dim \Delta_I \\ &= d(c_I - 1). \end{aligned}$$

Finally, observe that

$$Y_0 \mathcal{T} = \bigcap_{G \in \mathcal{T}} G = \bigcap_{I \in \mathcal{S}} \Delta_I = \Delta_{\mathcal{S}} \cong X^{c(\mathcal{S})}.$$

Therefore the expected conclusion is implied by Theorem 3.1. $\square$

Proof of Theorem 4.2. The statement of the motive decomposition is proved exactly as the above proof.

The correspondences are induced from Theorem 3.2. The improvement of this theorem than Theorem 3.2 is: we can say more about the Chern classes appeared in the correspondence $\alpha_{\mathcal{S}, \underline{\mu}}$ in Theorem 3.2.

First, given $G = \Delta_I$, let $\Pi = (I_1, \dots, I_{c_I})$ be the partition containing all sons of I in $\mathcal{S}$. We compute the normal bundle $N_G := N_{\Delta_I} \Delta_\Pi$. Without loss of generality, assume $I = (12 \dots m)$, where $m \leq n$.

Denote by $p_i : \Delta_I \rightarrow X$ and $q_i : \Delta_\Pi \rightarrow X$ the projections induced from the projection of X^n to the i -th factor. For each $1 \leq i \leq c_I$, pick an $a_i \in I_i$.

$$\begin{aligned} T_{\Delta_I} &= p_1^* T_X \oplus p_{m+1}^* T_X \oplus \dots \oplus p_n^* T_X, \\ T_{\Delta_\Pi} &= q_{a_1}^* T_X \oplus \dots \oplus q_{a_{c_I}}^* T_X \oplus q_{m+1}^* T_X \oplus \dots \oplus q_n^* T_X, \\ T_{\Delta_\Pi}|_{\Delta_I} &= p_1^* T_X \oplus \dots \oplus p_1^* T_X \oplus q_{m+1}^* T_X \oplus \dots \oplus q_n^* T_X, \end{aligned}$$

therefore, $c(N_G) = p_1^* c(T_X)^{c_I - 1}$.

To compute the Chern classes of N_G twisted by a line bundle L , we use the Chern root technique. For any vector bundle N on X , define the Chern polynomial as

$$c_y(N) := c_0(N) + c_1(N)y + c_2(N)y^2 + \dots$$

Define $x = c_1(L)$. Recall that the rank of N_G is $r_G = d(c_I - 1)$. Then

$$\begin{aligned} c(N_G \otimes L) &= c_{r_G}(N_G) + c_{r_G-1}(N_G)(1+x) + \dots + c_0(N_G)(1+x)^{r_G} \\ &= (x+1)^{r_G} c_{\frac{1}{x+1}}(N_G) \\ &= (x+1)^{d(c_I-1)} p_1^* c_{\frac{1}{x+1}}(T_X)^{c_I-1} \\ &= p_1^* [(x+1)^d c_{\frac{1}{x+1}}(T_X)]^{c_I-1} = p_1^* \zeta(x)^{c_I-1}. \end{aligned}$$

Finally, by restricting to $\Delta_{\mathcal{S}}$ and pulling back to $X(\mathcal{S})$ we get the expected formula for correspondences $\alpha_{\mathcal{S}, \underline{\mu}}$. $\square$

4.2. A formula for the generating function of Chow groups and Chow motive of $X[n]$. In this section, we express the decompositions of the Chow groups (Theorem 4.1) and the Chow motive (Theorem 4.2) in terms of exponential generating functions.

Define $[x^i]$ to be the function that picks up the coefficient of x^i from a power series. Define $[\frac{x^i t^n}{n!}]$ to be the function that picks up the coefficient of $\frac{x^i t^n}{n!}$ from a power series with two variables x and t , i.e.,

$$\left[\frac{x^i t^n}{n!} \right] \sum_{j,m} a_{jm} \frac{x^j t^m}{m!} := a_{in}.$$

The main theorem of this section is the following:

Theorem 4.3. *Define $f_i(x)$ to be the polynomials whose exponential generating function $N(x, t) = \sum_{i \geq 1} f_i(x) \frac{t^i}{i!}$ satisfies the identity*

$$(1-x)x^d t + (1-x^{d+1}) = \exp(x^d N) - x^{d+1} \exp(N).$$

Then for a nonsingular d -dimensional quasi-projective variety X ,

$$A^*(X[n]) = \bigoplus_{\substack{1 \leq k \leq n \\ i \geq 0}} A^{*-i}(X^k)^{\oplus [\frac{x^i t^n}{n!}] \frac{N^k}{k!}}.$$

Moreover, if X is projective, we have the motive decomposition

$$\begin{aligned} h(X[n]) &= \bigoplus_{\substack{\Pi=(I_1, \dots, I_k) \\ \text{partition of } [n]}} (h(\Delta_{\Pi})(i))^{\oplus [x^i](f_{|I_1|}(x) \dots f_{|I_k|}(x))} \\ &= \bigoplus_{\substack{1 \leq k \leq n \\ i \geq 0}} (h(X^k)(i))^{\oplus [\frac{x^i t^n}{n!}] \frac{N^k}{k!}}. \end{aligned}$$

Remark: One can write down by hand the first several terms of N . Define $\sigma_j = \sum_{i=1}^{dj-1} x^i$ (when $d = 1$, define $\sigma_1 = 0$). Then

$$\begin{aligned} N = & t + \sigma_1 \frac{t^2}{2!} + (\sigma_2 + 3\sigma_1^2) \frac{t^3}{3!} + (\sigma_3 + 10\sigma_1\sigma_2 + 15\sigma_1^3) \frac{t^4}{4!} \\ & + (\sigma_4 + 15\sigma_1\sigma_3 + 10\sigma_2^2 + 105\sigma_1^2\sigma_2 + 105\sigma_1^4) \frac{t^5}{5!} + \dots \end{aligned}$$

Proof of Theorem 4.3. We prove only the statement for motives, since the statement for Chow groups can be proved by exactly the same method.

By Theorem 4.2, we want to count for any given i and k , how many possible $\mathcal{S}$ and $\underline{\mu} \in \mathcal{S}$ satisfy $c(\mathcal{S}) = k$ and $\|\underline{\mu}\| = i$. First, consider the case when $c(\mathcal{S}) = 1$, i.e. $\mathcal{S}$ is a connected forest.

Define

$$f_n(x) := \sum_{\mathcal{S}: c(\mathcal{S})=1} \sum_{\underline{\mu} \in M_{\mathcal{S}}} x^{\|\underline{\mu}\|},$$

and define $f_1(x) = 1$.

For a nest $\mathcal{S}$ of $[n]$ with $c(\mathcal{S}) = 1$, we have

$$\sum_{\underline{\mu} \in M_{\mathcal{S}}} x^{\|\underline{\mu}\|} = \prod_{I \in \mathcal{S}^\circ} \sigma_{(c_I-1)},$$

i.e., I goes through all non-leaves of $\mathcal{S}$ (if $n = 1$, then the sum is set to be 1). Since the sons of the root of $\mathcal{S}$ correspond to a partition $\{I_1, \dots, I_k\}$ of $[n]$, we have following formula for $n \geq 2$,

$$f_n(x) = \sum_{\{I_1, \dots, I_k\} \text{ partition of } [n]} f_{|I_1|} f_{|I_2|} \dots f_{|I_k|} \sigma_{k-1}.$$

where $\sigma_k = \sum_{i=1}^{dk-1} x^i$ for $k > 0$, and $\sigma_0 = 0$. Since the equality does not hold for $n = 1$ where $f_1(x) = 1$ but the right side is 0, so one define

$$\tilde{f}_n(x) = \begin{cases} f_n(x), & \text{if } n > 1; \\ 0, & \text{if } n = 1. \end{cases}$$

Then the following holds for any $n \geq 1$:

$$\tilde{f}_n(x) = \sum_{\{I_1, \dots, I_k\} \text{ partition of } [n]} \tilde{f}_{|I_1|} \tilde{f}_{|I_2|} \dots \tilde{f}_{|I_k|} \sigma_{k-1}.$$

Recall the Compositional Formula of exponential generating functions (cf. [St99], Theorem 5.1.4), which asserts that if an equation as above holds, then

$$E_{\tilde{f}}(t) = E_{\sigma}(E_f(t)),$$

where

$$\begin{aligned} E_{\tilde{f}}(t) &= 1 + \tilde{f}_1 t + \tilde{f}_2 t^2/2! + \tilde{f}_3 t^3/3! + \dots \\ E_{\sigma}(t) &= 1 + \sigma_0 t + \sigma_1 t^2/2! + \sigma_2 t^3/3! + \dots \\ E_f(t) &= f_1 t + f_2 t^2/2! + f_3 t^3/3! + \dots \end{aligned}$$

By the definition of $\tilde{f}$, $E_{\tilde{f}} = E_f - t + 1$. Denote $N = E_f$, one has

$$N - t + 1 = E_g(N),$$

Standard Computation shows

$$E_g(N) = 1 + N + \frac{1}{x-1} \left[\frac{1}{x^d} (e^{x^d N} - 1) - x e^N + x \right].$$

Therefore

$$(1-x)x^d t + (1-x^{d+1}) = \exp(x^d N) - x^{d+1} \exp(N).$$

Now consider the case when $c(\mathcal{S})$ is not necessarily 1, i.e., the forest $\mathcal{S}$ is not necessarily connected. For a partition $\Pi = \{I_1, \dots, I_k\}$ of $[n]$, the number of times that $h(\Delta_\Pi)(i)$ appears in the decomposition of $h(X[n])$ is equal to $[x^k](f_{|I_1|}(x) \dots f_{|I_k|}(x))$, the coefficient of x^k in the product. Denote by $a_{k,i}$ the sum of these numbers for all partitions with k blocks. Then $a_{k,i}$ is the number of times that $h(X^k)(i)$ appears in the decomposition of $H(X[n])$.

Define

$$F_n(y) = \sum_{\{I_1, \dots, I_k\} \text{ partition of } [n]} f_{|I_1|} f_{|I_2|} \dots f_{|I_k|} y^k.$$

Then the coefficient $[y^k]F_n(y) = \sum a_{k,i} x^i$. Use the Compositional Formula again,

$$F_n = \left[\frac{t^n}{n!} \right] \exp(yN).$$

Therefore

$$\begin{aligned} [y^k]F_n(y) &= [y^k] \left[\frac{t^n}{n!} \right] \exp(yN) \\ &= \left[\frac{t^n}{n!} \right] [y^k] \exp(yN) \\ &= \left[\frac{t^n}{n!} \right] \frac{N^k}{k!}. \end{aligned}$$

This yields the formula for the decomposition of the Chow motive $h(X[n])$. $\square$

4.3. Description of $X[n]$ for small n . In this section we explain the previous Theorems (4.1, 4.2, and 4.3) about Fulton-MacPherson configuration space $X[n]$ for small $n = 2, 3, 4$.

For unification of expression, assume $d > 1$ in the following examples. (The case $d = 1$ is simpler but the expression needs to be modified.)

Example $n = 2$. The morphism $\pi : X[2] \rightarrow X^2$ is a blow-up along the diagonal Δ_{12} . Theorem 4.3 asserts

$$(8) \quad h(X[2]) \cong h(X^2) \oplus \bigoplus_{i=1}^{d-1} h(\Delta_{12})(i) \cong h(X^2) \oplus \bigoplus_{i=1}^{d-1} h(X)(i).$$

There are 2 possible nests: $\mathcal{S} = \{1, 2\}$ and $\mathcal{S} = \{1, 2, 12\}$. Theorem 4.2 asserts the following.

For the first nest, M_S contains only one element $\underline{\mu}$ with $\|\underline{\mu}\| = 0$. Therefore $\alpha = \Gamma_\pi$, $\beta = \Gamma_\pi^t$, $p = \Gamma_\pi^t \circ \Gamma_\pi$. They give the first direct summand in the decomposition (8).

For the second nest, $S^\circ = \{12\}$, $1 \leq \mu_{12} \leq d-1$, so there are $d-1$ direct summands for this nest. Denote $j : D_{12} \hookrightarrow X[2]$, $g : D_{12} \rightarrow \Delta_{12}$ as the natural map, we have

$$\begin{aligned}\alpha_{S,\underline{\mu}} &= -(j, g)_* j^* \left(\sum_{i=0}^{d-1-\mu_{12}} p_1^* c_i(T_X)(-D_{12})^{d-1-\mu_{12}-i} \right), \\ \beta_{S,\underline{\mu}} &= (g, j)_* j^* (D^{\mu_{12}-1}), \\ p_{S,\underline{\mu}} &= \beta_{S,\underline{\mu}} \circ \alpha_{S,\underline{\mu}}.\end{aligned}$$

They give the direct summand $h(\Delta_{12})(\mu_{12})$ in the decomposition (8).

Example $n = 3$. Apply Theorem 4.3,

$$\begin{aligned}h(X[3]) &\cong h(X^3) \oplus \bigoplus_{i=1}^{d-1} h(\Delta_{12})(i) \oplus \bigoplus_{i=1}^{d-1} h(\Delta_{13})(i) \oplus \bigoplus_{i=1}^{d-1} h(\Delta_{23})(i) \\ &\quad \oplus \bigoplus_{i=1}^{2d-1} (h(\Delta_{123})(i))^{\oplus \min\{3i-2, 6d-3i-2\}} \\ &\cong h(X^3) \oplus \bigoplus_{i=1}^{d-1} (h(X^2)(i))^{\oplus 3} \oplus \bigoplus_{i=1}^{2d-1} (h(X)(i))^{\oplus \min\{3i-2, 6d-3i-2\}}\end{aligned}$$

Now we write out all the correspondences that give the decomposition of motives. There are 8 possible nests, correspond to 8 trees (see the right side of Figure 4.3).

The tree on the left side of Figure 4.3 helps us to understand the relation between subvarieties of different Y_i 's (i.e. at different levels): each node with label I at level k correspond to the subvariety $Y_k I := (\Delta_I)^{(k)}$ in Y_k . The node at level k without label correspond to Y_k . For example, the root at level 4 corresponds to Y_4 , its two successors correspond to Y_3 and $Y_3(23)$, and the relation is that Y_4 is the blow-up of Y_3 along $Y_3(23)$.

We list below those correspondences α, β, p for the 8 trees:

① gives $\alpha = \Gamma_\pi, \beta = \Gamma_\pi^t, p = \Gamma_\pi^t \circ \Gamma_\pi$.




FIGURE 1. $X[3]$ by the symmetric construction.

② (and ③, ④ are similar) gives

$$\begin{aligned}\alpha_{\mathcal{S}, \underline{\mu}} &= (j_{\mathcal{S}}, g_{\mathcal{S}})_* j_{\mathcal{S}}^* \left(\{-p_1^* \zeta(-D_{123}) \frac{1}{1+D_{12}}\}_{d-1-\mu_{12}} \right), \\ \beta_{\mathcal{S}, \underline{\mu}} &= (g_{\mathcal{S}}, j_{\mathcal{S}})_* j_{\mathcal{S}}^* (D_{12}^{\mu_{12}-1}).\end{aligned}$$

where $X(\mathcal{S}) = D_{12}$, $1 \leq \mu_{12} \leq d-1$.

⑤ gives

$$\begin{aligned}\alpha_{\mathcal{S}, \underline{\mu}} &= (j_{\mathcal{S}}, g_{\mathcal{S}})_* j_{\mathcal{S}}^* \left(\{-p_1^* \zeta(\mathcal{O})^2 \frac{1}{1+D_{123}}\}_{2d-1-\mu_{123}} \right), \\ \beta_{\mathcal{S}, \underline{\mu}} &= (g_{\mathcal{S}}, j_{\mathcal{S}})_* j_{\mathcal{S}}^* (D_{123}^{\mu_{123}-1}).\end{aligned}$$

where $X(\mathcal{S}) = D_{123}$, $1 \leq \mu_{123} \leq 2d-1$.

⑥ (and ⑦, ⑧ are similar) gives

$$\begin{aligned}\alpha_{\mathcal{S}, \underline{\mu}} &= (j_{\mathcal{S}}, g_{\mathcal{S}})_* j_{\mathcal{S}}^* \\ &\quad \left(\{p_1^* \zeta(-D_{123}) \frac{1}{1+D_{12}}\}_{d-1-\mu_{12}} \{p_1^* \zeta(\mathcal{O}) \frac{1}{1+D_{123}}\}_{d-1-\mu_{123}} \right), \\ \beta_{\mathcal{S}, \underline{\mu}} &= (g_{\mathcal{S}}, j_{\mathcal{S}})_* j_{\mathcal{S}}^* (D_{12}^{\mu_{12}-1} D_{123}^{\mu_{123}-1}).\end{aligned}$$

where $X(\mathcal{S}) = D_{12} \cap D_{123}$, $1 \leq \mu_{12}, \mu_{123} \leq d-1$.

Remark: If we use Fulton and MacPherson's nonsymmetric construction of $X[3]$, we would get another set of correspondences which also gives a decomposition of the motive $h(X[n])$. This set of correspondences turns out to be different than the ones given above: a straightforward calculation shows that, by the nonsymmetric construction of $X[3]$, the correspondence that gives the direct summand $h(\Delta_{12})(\mu_{12})$ is


$$\begin{aligned}\alpha : \quad h(X[3]) &\rightarrow h(\Delta_{12})(\mu_{12}), \\ \alpha &= (j_{12}, g_{12})_* j_{12}^* \left(\{p_1^* \zeta(\mathcal{O}) \frac{1}{1+D_{12}}\}_{d-1-\mu_{12}} \right).\end{aligned}$$

where $j_{12} : D_{12} \hookrightarrow X[3]$ and $g_{12} : D_{12} \rightarrow \Delta_{12}$ are the natural morphisms. However, the correspondence giving the direct summand $h(\Delta_{13})(\mu_{13})$ is

$$\begin{aligned}\alpha' : \quad h(X[3]) &\rightarrow h(\Delta_{13}) \otimes \mathbb{L}^{\mu_{13}}, \\ \alpha' &= (j_{13}, g_{13})_* j_{13}^* \left(\{p_1^* \zeta(-D_{123}) \frac{1}{1+D_{13}}\}_{d-1-\mu_{13}} \right).\end{aligned}$$

where $j_{13} : D_{13} \hookrightarrow X[3]$, $g_{13} : D_{13} \rightarrow \Delta_{13}$ are the natural morphisms. Notice that α and α' are not of similar forms (Compare $\zeta(\mathcal{O})$ with $\zeta(-D_{123})$). Therefore the non-symmetry of the construction of $X[3]$ induces the non-symmetry of correspondences. Actually, this is one reason why we choose the symmetric construction of $X[n]$ (cf. Remark 4.1).

Example $n = 4$. we only look at one nest $\mathcal{S}$:



We have $X(\mathcal{S}) = D_{12} \cap D_{34}$, $1 \leq \mu_{12}, \mu_{34} \leq d-1$ and

$$\begin{aligned} \alpha_{\mathcal{S}, \underline{\mu}} &= (j_{\mathcal{S}}, g_{\mathcal{S}})_* j_{\mathcal{S}}^* \\ &\quad \left(\{p_1^* \zeta(-D_{1234}) \frac{1}{1+D_{12}}\}_{d-1-\mu_{12}} \{p_3^* \zeta(-D_{1234}) \frac{1}{1+D_{34}}\}_{d-1-\mu_{34}} \right), \\ \beta_{\mathcal{S}, \underline{\mu}} &= (g_{\mathcal{S}}, j_{\mathcal{S}})_* j_{\mathcal{S}}^* (D_{12}^{\mu_{12}-1} D_{34}^{\mu_{34}-1}). \end{aligned}$$

Since Δ_{12} and Δ_{34} would not be disjoint in the procedure of blow-ups, a priori we have to make a choice of order that whether blow up along (the strict transform of) Δ_{12} first, or along (the strict transform of) Δ_{34} first. Although we have to choose (non-canonically) an order to compute the correspondences, it turns out that the correspondences (hence projectors) which give the motive decomposition in Theorem 4.2 are actually independent of the choice, therefore “canonical”. This independence is a special case of Remark 4.1: for $\sigma = (13)(24) \in \mathbb{S}_4$, the above correspondences is invariant under the action induced by σ .

An application of Theorem 4.3 is: we can compute the rank of $A(X[n])$ (as an abelian group) once given the ranks of $A(X^k)$ for all $1 \leq k \leq n$ (assuming that the ranks of $A(X^k)$ ’s are finite).

Let us take $\mathbb{P}^d[5]$ for example. Since the rank of $A((\mathbb{P}^d)^k)$ is $(d+1)^k$, Theorem 4.3 implies that the rank of $A(\mathbb{P}^d[5])$ is

$$\sum_{1 \leq k \leq 5} (d+1)^k \left(\left[\frac{t^5}{t!} \right] \left(\frac{N^k}{k!} \Big|_{x=1} \right) \right).$$

By Remark 4.2, we can compute the following

$$\begin{aligned} \frac{N^2}{2!} &= \frac{t^2}{2!} + 3\sigma_1 \frac{t^3}{3!} + (15\sigma_1^2 + 4\sigma_2) \frac{t^4}{4!} + (105\sigma_1^3 + 60\sigma_1\sigma_2 + 5\sigma_3) \frac{t^5}{5!} + \dots \\ \frac{N^3}{3!} &= \frac{t^3}{3!} + 6\sigma_1 \frac{t^4}{4!} + (45\sigma_1^2 + 10\sigma_2) \frac{t^5}{5!} + \dots \\ \frac{N^4}{4!} &= \frac{t^4}{4!} + 10\sigma_1 \frac{t^5}{5!} + \dots \\ \frac{N^5}{5!} &= \frac{t^5}{5!} + \dots \end{aligned}$$

Now plug in $x = 1$, we have $\sigma_j = dj - 1$. The above sum is a polynomial of d as follows

$$\begin{aligned} &(d+1)^5 + (d+1)^4 10\sigma_1 + (d+1)^3 (45\sigma_1^2 + 10\sigma_2) \\ &+ (d+1)^2 (105\sigma_1^3 + 60\sigma_1\sigma_2 + 5\sigma_3) \\ &+ (d+1) (\sigma_4 + 15\sigma_1\sigma_3 + 10\sigma_2^2 + 105\sigma_1^2\sigma_2 + 105\sigma_1^4). \end{aligned}$$

In particular, the rank of $A(\mathbb{P}^1[5])$ is 178, the rank of $A(\mathbb{P}^2[5])$ is 7644.

Remark: For the example $X = \mathbb{P}^d$, since $X[n]$ has an affine cell decomposition, the rank of the Chow group $A_k(X[n])$ coincides with the $2k$ -th Betti number of $X[n]$. Therefore we could also get the above rank by the Poincaré polynomial of $X[n]$ computed in [FM94]. However, the rank of $A(X[n])$ for a general variety X is not implied by the Poincaré polynomial of $X[n]$.

5. CHOW MOTIVES OF $X[n]/\mathfrak{S}_n$

It is proved in [FM94] that the isotropy group of any point in $X[n]$ is a solvable group. It is natural to consider the quotient space $X[n]/\mathfrak{S}_n$. In this section, we compute its Chow motive in terms of the Chow motives of the Cartesian products of symmetric products of X .

The base field is of characteristic 0 throughout this section.

Lemma 5.1. *Suppose a finite group G acts on a nonsingular projective variety Y . If $p_1, \dots, p_k$ are orthogonal projectors of Y that*

i) $\sigma p_i = p_i \sigma, \forall 1 \leq i \leq k, \forall \sigma \in G$.

ii) $p_1 + p_2 + \dots + p_k = \Delta_Y$.

Then $\text{ave} \Delta_Y = \sum \text{ave} \circ p_i$ where $\text{ave} \circ p_1, \dots, \text{ave} \circ p_k$ are orthogonal projectors. Consequently, $h(Y) = \oplus (Y, \text{ave} \circ p_i)$.

Proof. Since

$$\begin{aligned} (\text{ave } p_i)(\text{ave } p_j) &= \left(\frac{1}{|G|} \sum_{\sigma} \sigma p_i \right) \left(\frac{1}{|G|} \sum_{\tau} \tau p_j \right) \\ &= \frac{1}{|G|^2} \sum_{\sigma, \tau} \sigma \tau p_i p_j = \frac{1}{|G|} \sum_{\sigma} \sigma \delta_{ij} p_i = \delta_{ij} (\text{ave } p_j). \end{aligned}$$

Then the lemma follows. $\square$

Lemma 5.2. *Suppose Y, Z are nonsingular (not necessary connected) projective varieties with finite group G actions. Suppose that $\alpha \in \text{Corr}^{-m}(Y, Z)$ has an inverse $\beta \in \text{Corr}^m(Z, Y)$, and α gives an isomorphism of Chow motives*

$$(Y, p) \cong h(Z)(m)$$

where $p = \beta \alpha$, and $\alpha \sigma = \sigma \alpha, \beta \sigma = \sigma \beta, \forall \sigma \in G$. Then

$$(Y, \text{ave} \circ p) \cong h(Z/G)(m).$$

Proof. Similar to the proof of Lemma 5.1, we have $(\text{ave } p)^2 = \text{ave } p$ and the following commutative diagram

$$\begin{array}{ccccc} Y & \xrightarrow{\text{ave } \alpha} & Z & \xrightarrow{\text{ave } \beta} & Y \\ \text{ave } p \downarrow & & \downarrow \text{ave } \Delta_Z & & \downarrow \text{ave } p \\ Y & \xrightarrow{\text{ave } \alpha} & Z & \xrightarrow{\text{ave } \beta} & Y. \end{array}$$

Therefore, $(Y, \text{ave } p) \cong (Z, \text{ave } \Delta_Z)(m) \cong h(Z/G)(m)$. $\square$

Now we consider the quotient variety $X[n]/\mathfrak{S}_n$. For convenience, define $G := \mathfrak{S}_n$. There is a natural action of G on the set $\{(\mathcal{S}, \underline{\mu})\}$ where $\mathcal{S}$ are nests and $\underline{\mu} \in M_{\mathcal{S}}$. Define the subgroup $G_{\mathcal{S}, \underline{\mu}}$ of G as

$$G_{\mathcal{S}, \underline{\mu}} = \{\sigma \in \mathfrak{S}_n : \sigma(\mathcal{S}, \underline{\mu}) = (\mathcal{S}, \underline{\mu})\}.$$

Define $\overline{(\mathcal{S}, \underline{\mu})}$ to be the class of G -orbit $G \cdot (\mathcal{S}, \underline{\mu})$. Then

$$\Delta_Y = \sum_{\mathcal{S}, \underline{\mu}} p_{\mathcal{S}, \underline{\mu}} = \sum_{\overline{(\mathcal{S}, \underline{\mu})}} \sum_{\bar{\sigma} \in G/G_{\mathcal{S}, \underline{\mu}}} p_{\sigma(\mathcal{S}, \underline{\mu})}.$$

Since $\{\alpha_{\mathcal{S}, \underline{\mu}}\}, \{\beta_{\mathcal{S}, \underline{\mu}}\}$ are $\mathfrak{S}_n$ -symmetric (cf. the Remark after Theorem 4.2), it is easy to check that $\sum_{\bar{\sigma} \in G/G_{\mathcal{S}, \underline{\mu}}} p_{\sigma(\mathcal{S}, \underline{\mu})}$ commutes with every $\tau \in G$. By Lemma 5.1,

$$h(X[n]/G) \cong (Y, \text{ave} \circ \Delta_Y) \cong \bigoplus_{\overline{(\mathcal{S}, \underline{\mu})}} \left(Y, \text{ave} \circ \sum_{\bar{\sigma} \in G/G_{\mathcal{S}, \underline{\mu}}} p_{\sigma(\mathcal{S}, \underline{\mu})} \right).$$

Since

$$\left(Y, \sum_{\bar{\sigma} \in G/G_{\mathcal{S}, \underline{\mu}}} p_{\sigma(\mathcal{S}, \underline{\mu})} \right) \cong \left(\bigsqcup_{\bar{\sigma} \in G/G_{\mathcal{S}, \underline{\mu}}} \Delta_{\sigma(\mathcal{S})} \right) (||\underline{\mu}||),$$

by Lemma 5.2 we have

$$\begin{aligned} \left(Y, \text{ave} \circ \sum_{\bar{\sigma} \in G/G_{\mathcal{S}, \underline{\mu}}} p_{\sigma(\mathcal{S}, \underline{\mu})} \right) &\cong h \left(\left(\bigsqcup_{\bar{\sigma} \in G/G_{\mathcal{S}, \underline{\mu}}} \Delta_{\sigma(\mathcal{S})} \right) / G \right) (||\underline{\mu}||) \\ &\cong h(\Delta_{\mathcal{S}}/G_{\mathcal{S}, \underline{\mu}}) (||\underline{\mu}||). \end{aligned}$$

The space $\Delta_{\mathcal{S}}/G_{\mathcal{S}, \underline{\mu}}$ can be described as follows. Each $(\mathcal{S}, \underline{\mu})$ corresponds to a labeled “weighted” forest, the correspondence is given by attaching an integer μ_I to each non-leaf node I of the labeled forest $\mathcal{S}$. Forgetting all the labels on the nodes of $\mathcal{S}$, we get an unlabeled weighted forest of the form $n_1 T_1 + \dots + n_r T_r$, where T_i are mutually distinct unlabeled weighted tree (we call such a tree is of type $\{n_1, \dots, n_r\}$). Then

$$\Delta_{\mathcal{S}}/G_{\mathcal{S}, \underline{\mu}} \cong X^{(n_1)} \times \dots \times X^{(n_r)}$$

Figure 2 gives an example of a labeled weighted forest and the corresponding unlabeled weighted forest. The weight a, b are integers.



FIGURE 2. Labeled and unlabeled weighted forests.

Therefore we have proved the following decomposition of the Chow motive of $X[n]/\mathfrak{S}_n$:

Theorem 5.3. *For any unordered set of integers $\nu = \{n_1, \dots, n_r\}$ and any integer m , let $\lambda(\nu, m)$ to be the number of unlabeled weighted forest with n leaves, of type ν and total weight m , such that at each non-leaf v with c_v children, the weight m_v satisfies $1 \leq m_v \leq (c_v - 1) \dim X - 1$. Then*

$$h(X[n]/\mathfrak{S}_n) = \bigoplus_{\nu, m} \left[h(X^{(n_1)} \times \dots \times X^{(n_r)})(m) \right]^{\oplus \lambda(\nu, m)}.$$

Remark: An application of this theorem. MacDonald proved a formula that relates the Betti number of X and its symmetric powers:

$$\sum_{n=0}^{\infty} P_t X^{(n)} \cdot T^n = \frac{(1+tT)^{b_1} (1+t^3T)^{b_3} \dots}{(1-T)^{b_0} (1-t^2T)^{b_2} \dots}$$

where b_i is the i -th Betti number of X . By the decomposition of the de Rham cohomology of $X[n]/\mathfrak{S}_n$ induced by the motivic decomposition formula in the above theorem, we can compute the Betti number of $X[n]/\mathfrak{S}_n$ (modulo the combinatorial difficulty of calculating $\lambda(\nu, m)$).

Examples: Here are some examples of $h(X[n]/\mathfrak{S}_n)$ for small n . Let $d = \dim X$.


i) $n=2$. There are d different forests as follows, where each weight $a \in \mathbb{Z}$ ($1 \leq a \leq d-1$) gives a forest:



Therefore

$$h(X[2]/\mathfrak{S}_2) \cong h(X^{(2)}) \oplus \bigoplus_{a=1}^{d-1} h(X)(a).$$

ii) $n=3$. The forests are:



where the weights $a, b, c, e \in \mathbb{Z}$ satisfy $1 \leq a, c, e \leq d-1$, and $1 \leq b \leq 2d-1$. We have

$$h(X[3]/\mathfrak{S}_3) \cong h(X^{(3)}) \oplus \bigoplus_{i=1}^{d-1} (h(X^2)(i))^{\oplus 3} \oplus \bigoplus_{i=1}^{2d-1} (h(X)(i))^{\oplus \min\{i, 2d-i\}}.$$

iii) $n=4$. The varieties appear in the decomposition of $h(X[4]/\mathfrak{S}_4)$ are:

$$X^{(4)}, X \times X^{(2)}, X^2, X^{(2)}, X.$$

The decomposition is a bit nasty to be written here. Therefore we only point out a fact. Consider the forest in Figure 3, where $a, b \in \mathbb{Z}$ and $1 \leq a, b \leq d-1$. For any $a < b$, the weighted forest is of type $\nu = \{1, 1\}$ and therefore



FIGURE 3. An unlabeled weighted forest when $n = 4$

gives a summand $h(X^2)(a+b)$. However, for $a = b$, this weighted forest has an automorphism exchanging the two trees, thus is of type $\nu = \{2\}$ and gives a summand $h(X^{(2)})(2a)$. Due to this kind of automorphism of weighted forests, it seems difficult to compute $\lambda(\nu, m)$.

Question. *Is there a clean formula for $\lambda(\nu, m)$? (Maybe in terms of a generating function?)*

APPENDIX A. A FORMULA FOR THE MOTIVE OF A BLOW-UP

Suppose $f : \tilde{Y} \rightarrow Y$ is the blow-up of a nonsingular projective variety Y along a nonsingular closed subvariety V of Y , and denote by P the exceptional divisor. Denote by i, j, f, g the morphisms as in the following fibre square

$$\begin{array}{ccc} P & \xrightarrow{j} & \tilde{Y} \\ g \downarrow & \square & \downarrow f \\ V & \xrightarrow{i} & Y. \end{array}$$

Denote by $N := N_V Y$ the normal bundle of V in Y . Let $h := c_1(\mathcal{O}_N(1)) \in A^1(P)$. Let $r := \text{codim}_V Y$ be the codimension of V in Y .

For $1 \leq k \leq r-1$, define $\alpha_k \in \text{Corr}^{-k}(\tilde{Y}, V)$, $\beta_k \in \text{Corr}^k(V, \tilde{Y})$, $p_k \in \text{Corr}^0(\tilde{Y}, \tilde{Y})$, $\alpha_0 \in \text{Corr}^0(\tilde{Y}, Y)$, $\beta_0 \in \text{Corr}^0(Y, \tilde{Y})$ and $p_0 \in \text{Corr}^0(\tilde{Y}, \tilde{Y})$ as follows

$$(9) \quad \left\{ \begin{array}{l} \alpha_0 := \Gamma_f, \\ \beta_0 := \Gamma_f^t, \\ p_0 := \beta_0 \circ \alpha_0 = \Gamma_f^t \circ \Gamma_f = (f \times f)^* \Delta_Y, \\ \alpha_k := -(j, g)_* \left(\sum_{l=0}^{r-1-k} g^* c_{r-1-k-l}(N) h^l \right) \\ \quad = -(j, g)_* \left(\left\{ g^* c(N) \frac{1}{1-h} \right\}_{r-1-k} \right), \\ \beta_k := (g, j)_* h^{k-1}, \\ p_k := \beta_k \circ \alpha_k, \end{array} \right.$$

where the subscript $r - 1 - k$ in the definition of α_k means taking the codimension $(r - 1 - k)$ component. We will give the proof of the following proposition at the end of this section.

Proposition A.1. *Define $\alpha_k, \beta_k, p_k, \alpha_0, \beta_0, p_0$ as above. The following holds.*

(i) $\alpha_0\beta_0 = \Delta_Y$, $\alpha_k\beta_k = \Delta_V$ for $1 \leq k \leq r - 1$;

$\alpha_i\beta_j = 0$ for $i \neq j$.

(ii) $p_0, p_1, p_2, \dots, p_{r-1}$ are mutually orthogonal projectors of $\tilde{Y}$, and

$$\sum_{i=0}^{r-1} p_i = \Delta_{\tilde{Y}} \quad \text{in } A(\tilde{Y} \times \tilde{Y}),$$

i.e. equality holds up to rational equivalence.

(iii) We have the following isomorphisms of motives,

$\alpha_0 : (\tilde{Y}, p_0, 0) \simeq h(Y)$, with inverse morphism β_0 ,

$\alpha_k : (\tilde{Y}, p_k, 0) \simeq h(V)(k)$, with inverse morphism β_k , for $1 \leq k \leq r - 1$.

Define $\Gamma := \bigoplus_{i=0}^{r-1} \alpha_i$, $\Gamma' := \sum_{i=0}^{r-1} \beta_i$, then Proposition A.1 can be conveniently reformulated as follows:

Theorem A.2. *The correspondence Γ gives a canonical isomorphism in CHM ,*

$$\Gamma : h(\tilde{Y}) \cong h(Y) \oplus \bigoplus_{k=1}^{r-1} h(V)(k).$$

with an inverse isomorphism given by Γ' .

Remark: When the normal bundle N of V in Y is trivial (for example, when V is a point), P is isomorphic to a product space $V \times \mathbb{P}^{r-1}$ and $h = c_1(\mathcal{O}_P(1))$ can be represented (not canonically) by a product space $H = V \times \mathbb{P}^{r-2}$ in P . In this case, we have simple forms for the projectors:

$$p_k = -(j \times j)_*(H^{r-1-k} \times_V H^{k-1}), \quad \text{for } 1 \leq k \leq r - 1;$$

$$p_0 = \Delta + \sum_{k=1}^{r-1} (j \times j)_*(H^{r-1-k} \times_V H^{k-1}).$$

In general, for a nontrivial normal bundle N , more terms involving the Chern classes of N are needed, and the correspondences cannot be represented by explicit and natural algebraic cycles.

Remark: The isomorphism of motives in Theorem A.2 is also a consequence of “Theorem on the additive structure of the motif” of $\tilde{Y}$ in [Man68] §9, which states, in our notation, that there is a split exact sequence

$$0 \longrightarrow h(V)(r) \xrightarrow{a} h(Y) \oplus h(P)(1) \xrightarrow{b} h(\tilde{Y}) \longrightarrow 0.$$

The correspondences appeared in our theorem are not given, at least not explicitly, in Manin's paper.

In order to clarify this point, define

$$\begin{aligned}\Phi &= c_{r-1}(g^*N/\mathcal{O}_N(-1)) \in A^{r-1}(P), c_\Phi = \delta_{P*}(\Phi) \in \text{Corr}(P, P), \\ a &= (i_*, c_\Phi \circ g^*), a' = g_*, \\ b &= f^* + j_*, \quad b' \text{ its right inverse,} \\ d &= \Delta_{Y \times P} - aa', d' = \Delta_Y \otimes (\Delta_P - p_0^P) \text{ (where } p_0^P = c_{h^{r-1}} \circ g^* \circ g_*),\end{aligned}$$

denote by $e : \bigoplus_{k=1}^{r-1} V(k) \rightarrow (P, \Delta_P - p_0^P)$ the isomorphism implicitly defined in [Man68] §7, and denote by e' the inverse of e .

We have the following isomorphisms

$$\begin{aligned}h(Y) \oplus \bigoplus_{k=1}^{r-1} h(V)(k) &\xrightleftharpoons[\Delta_Y \otimes e']{\Delta_Y \otimes e} (Y \sqcup P, (\Delta_Y, \Delta_P - p_0^P)) \xrightleftharpoons[d']{d} \\ &(Y \sqcup P, \Delta_{Y \sqcup P} - aa') \xrightleftharpoons[b']{b} (\tilde{Y}, \Delta_{\tilde{Y}}).\end{aligned}$$

Hence the following is an isomorphism of Chow motives

$$(\Delta_Y \otimes e') \circ d' \circ b' : h(\tilde{Y}) \cong h(Y) \oplus \bigoplus_{k=1}^{r-1} h(V) \otimes \mathbb{L}^k$$

with inverse $b \circ d \circ (\Delta_Y \otimes e)$.

Therefore, to write down the correspondence $(\Delta_Y \otimes e') \circ d' \circ b'$, we need to find explicitly the right inverse b' of b . However in [Man68] the construction of b' is based on the surjectivity of $\gamma : A(\tilde{Y} \times (Y \sqcup P)) \rightarrow A(\tilde{Y} \times \tilde{Y})$ as follows: by the surjectivity of γ , there is a cycle class $c \in A(\tilde{Y} \times (Y \sqcup P))$ (which is not given, at least explicitly, in [Man68]) such that $\gamma(c) = \Delta_{\tilde{Y}} \in A(\tilde{Y} \times \tilde{Y})$. Then b' is defined to be $(1 - aa')c$.

On the other hand, the correspondences Γ and Γ' we have constructed in Theorem A.2 give an explicit construction of b' . Indeed, $b' = d \circ (\Delta_Y \otimes e) \circ \Gamma$.

Proof of Proposition A.1. In the proof, we assume $1 \leq k \leq r-1$, $0 \leq i, j \leq r-1$.

The idea is as follows: we study the morphisms α_{i*}, β_{i*} and p_{i*} of Chow groups induced by the correspondences α_i, β_i and p_i . As a consequence, the identities of morphisms of Chow groups which are induced by the identities in Proposition A.1 (i) (ii) hold. On the other hand, Manin's Identity Principle asserts that the identities of morphisms of Chow groups imply the identities of correspondences, providing that the correspondences are universal in certain sense.

By [Vo03] Theorem 9.27, an element $\tilde{y} \in A(\tilde{Y})$ can be expressed uniquely as

$$\tilde{y} = \sum_{i=1}^{r-1} j_*(g^*a_i \cdot h^{i-1}) + f^*y.$$

It is standard to verify

- (α_k) The morphism $\alpha_{k*} : A(\tilde{Y}) \rightarrow A(V)$ maps $\tilde{y} \mapsto a_k$.
- (β_k) The morphism $\beta_{k*} : A(V) \rightarrow A(\tilde{Y})$ maps $x \mapsto j_*(g^*x \cdot h^{k-1})$.
- (α_0) The morphism $\alpha_{0*} : A(\tilde{Y}) \rightarrow A(Y)$ maps $\tilde{y} \mapsto y$.
- (β_0) The morphism $\beta_{0*} : A(Y) \rightarrow A(\tilde{Y})$ maps $y \mapsto f^*y$.

To give a flavor, we prove only the statement (α_k), that is, $\alpha_{k*}(\tilde{y}) = a_k$. Define $a_0 = -i^*y$. Since $j^*j_*z = -h \cdot z$ for $\forall z \in A(P)$, we have

$$j^*\tilde{y} = \sum_{i=1}^{r-1} j^*j_*(g^*a_i \cdot h^{i-1}) + j^*f^*y = -\sum_{i=1}^{r-1} g^*a_i \cdot h^i + g^*i^*y = -\sum_{i=0}^{r-1} g^*a_i \cdot h^i.$$

By definition (see [Fu98] §3), the i -th Segre class of N is

$$s_i(N) := g_*(h^{i+r-1}),$$

hence

$$\begin{aligned} \alpha_{k*}(\tilde{y}) &= -g_*(j^*\tilde{y} \cdot \sum_{l=0}^{r-1-k} g^*c_{r-1-k-l}(N) \cdot h^l) \\ &= -g_*\left(\left(-\sum_{i=0}^{r-1} g^*a_i \cdot h^i\right) \cdot \left(\sum_{l=0}^{r-1-k} g^*c_{r-1-k-l} \cdot h^l\right)\right) \\ &= g_*\left(\sum_{i=0}^{r-1} \sum_{l=0}^{r-1-k} g^*(a_i c_{r-1-k-l}) h^{i+l}\right) \\ &= \sum_{i=0}^{r-1} a_i \left(\sum_{l=0}^{r-1-k} c_{r-1-k-l} s_{i+l+1-r}\right). \end{aligned}$$

Since $c(N)s(N) = 1$ where $c(N) := \sum c_i(N)$ is the total Chern class and $s(N) := \sum s_i(N)$ is the total Segre class, we have

$$\sum_{l=0}^{r-1-k} c_{r-1-k-l} s_{i+l+1-r} = \sum_{l=-\infty}^{+\infty} c_{r-1-k-l} s_{i+l+1-r} = \{c(N)s(N)\}_{i-k} = \delta_{ik},$$

the first equality is because $s_{i+l+1-r} = 0$ for $l < 0$ and $c_{r-1-k-l} = 0$ for $l > r-1-k$. It follows that $\alpha_{k*}(\tilde{y}) = a_k$, as we claimed.

The statements $(\alpha_k), (\beta_k), (\alpha_0), (\beta_0)$ immediately imply the following identities:

$$\begin{aligned} \alpha_{k*}\beta_{k*} &= id_{A(V)}, & \alpha_{0*}\beta_{0*} &= id_{A(Y)}, & \alpha_{i*}\beta_{j*} &= 0 \text{ for } i \neq j, \\ (p_i p_j)_* &= \delta_{ij} p_{i*}, & \sum_{i=0}^{r-1} p_{i*} &= id_{A(\tilde{Y})}. \end{aligned}$$

For any smooth scheme T , $T \times \tilde{Y}$ is the blow-up of $T \times Y$ along the smooth subvariety $T \times V$. Denote $j' = id_T \times j$, $g' = id_T \times g$, $f' = id_T \times f$, $i' = id_T \times i$, we have the following fiber square:

$$\begin{array}{ccc} T \times P & \xrightarrow{j'} & T \times \tilde{Y} \\ \downarrow g' & \square & \downarrow f' \\ T \times V & \xrightarrow{i'} & T \times Y \end{array}$$

We can construct the correspondences $\alpha'_i, \beta'_i, p'_i$ for this fiber square as we did in (9). we have

$$\alpha'_i = id_T \otimes \alpha_i, \beta'_i = id_T \otimes \beta_i, p'_i = id_T \otimes p_i.$$

Then (i) and (ii) follows from Manin's Identity Principle.

For (iii), to show that α_k gives an isomorphism $(\tilde{Y}, p_k, 0) \simeq h(V) \otimes \mathbb{L}^k$ with inverse β_k , we need to show that $p_k = p_k \circ \beta_k \circ \alpha_k$ and $id = id \circ \alpha_k \circ \beta_k$. but they are direct consequences of the fact that $\alpha_k \circ \beta_k = \Delta_V$ from (i). The proof for $(\tilde{Y}, p_0, 0) \simeq h(Y)$ is similar. $\square$

REFERENCES

- [AM69] M. Atiyah, I. Macdonald, Introduction to commutative algebra, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont. 1969 ix+128 pp.
- [CH00] A. Corti, M. Hanamura, *Motivic decomposition and intersection Chow groups. I*, Duke Math. J. 103 (2000), no. 3, 459–522.
- [dBV98] S. del Baño, V. Navarro, *On the motive of a quotient variety*, Dedicated to the memory of Fernando Serrano. Collect. Math. 49 (1998), no. 2-3, 203–226.
- [DP95] C. De Concini, C. Procesi, *Wonderful models of subspace arrangements*, Selecta Mathematica 1 (1995), 459–494.
- [Fu98] W. Fulton, Intersection theory, Second edition, Springer-Verlag, Berlin, 1998.
- [FM94] W. Fulton, R. MacPherson, *A compactification of configuration spaces*, Ann. Math. **139** (1994), 183–225.
- [Ha77] R. Hartshorne, Algebraic geometry, Graduate Texts in Mathematics, No. 52, Springer-Verlag, New York-Heidelberg, 1977.
- [Hu03] Y. Hu, *A compactification of open varieties*, Trans. Amer. Math. Soc. 355 (2003), no. 12, 4737–4753.
- [Li06] L. Li, *Wonderful compactifications of arrangements of subvarieties*, preprint math.AG/0611412, to appear in Michigan Math. J..
- [Ka93] M. Kapranov, *Chow quotients of Grassmannians*, I. I. M. Gel'fand Seminar, 29–110, Adv. Soviet Math., 16, Part 2, Amer. Math. Soc., Providence, RI, 1993.

- [Ke92] S. Keel, *Intersection theory of moduli space of stable n -pointed curves of genus zero*, Trans. Amer. Math. Soc. 330 (1992), no. 2, 545–574.
- [Ke93] S. Keel, *Intersection theory of linear embeddings* Trans. Amer. Math. Soc. 335 (1993), no. 1, 195–212.
- [KT99] G. Kuperberg, D. Thurston, *Perturbative 3-manifold invariants by cut-and-paste topology*, math.GT/9912167.
- [Man68] J.I. Manin, *Correspondences, motifs and monoidal transformations*, Math. USSR-Sbornik 6 (1968), 439–470.
- [MP98] R. MacPherson, C. Procesi, *Making conical compactifications wonderful*, Selecta Math. (N.S.) 4 (1998), no. 1, 125–139.
- [St99] R.P. Stanley, *Enumerative combinatorics*, Vol. 2., Cambridge Studies in Advanced Mathematics, 62. Cambridge University Press, Cambridge, 1999.
- [Th99] D. Thurston, *Integral Expressions for the Vassiliev Knot Invariants*, math.AG/9901110.
- [UI02] A. Ulyanov, *Polydiagonal compactification of configuration spaces*, J. Algebraic Geom. 11 (2002), no. 1, 129–159.
- [Vo03] C. Voisin, *Hodge theory and complex algebraic geometry. II*. Translated from the French by Leila Schneps. Cambridge Studies in Advanced Mathematics, 77. Cambridge University Press, Cambridge, 2003.

LI LI

Department of Mathematics

University of Illinois at Urbana-Champaign

Email: llpku@math.uiuc.edu